

Zool. 498 xe

THE
CONCHOLOGICAL
ILLUSTRATIONS,

BY

G. B. SOWERBY, JUN.


LONDON:

SOWERBY, 50, GREAT RUSSELL STREET,
BLOOMSBURY.

1841.

41 = D.

fol. 498 ^{xe}
—



GEORGE ODELL, PRINTER, PRINCES STREET, CAVENDISH SQUARE.

2⁶

NOTICE.

THE artist and proprietor feels unwilling to present this series of Illustrations to the scientific public, without taking advantage of the opportunity afforded by its completion, to acknowledge his obligations for the support he has received. During the periodical issue of the parts, he has experienced the utmost kindness and attention from collectors, who have materially assisted him by throwing open their cabinets for his inspection, and by freely allowing him the use of their new and rare shells. He has also enjoyed a degree of patronage in the sale of the parts, which, although not amounting to a remuneration for the labour, has sufficed to meet the pecuniary expenses attending their production, and also to afford the prospect of ultimate advantage. The 1,500 figures contained in the work, represent 1,000 new species and varieties of 21 different Genera, and include all the species of *Cypræa*, *Chilina*, *Neritina*, *Margarita*, *Eulima*, and *Typhis*.

The catalogues embrace 19 Genera, and contain a reference to the description and to the best figure of each species, as far as could be known, up to the date of their publication. Those which were published in the earlier part of the work are from the pen of Mr. G. B. Sowerby, Sen. For the four last, viz.—those of *Chiton*, *Cardium*, *Murex*, and *Typhis*, a younger and less experienced conchologist must be held responsible. With regard to the whole of them, it is hoped that although some errors and imperfections must have occurred, they will yet be found to present as complete and perfect accounts of the Genera of which they treat as could be expected, considering the intricacies and difficulties of the

task; and they will be found to assist materially in the advancement of conchological knowledge. The three remaining Genera, of which illustrations, but not catalogues, have been given, are accompanied by indices referring to descriptions and stating localities in every practicable instance.

A short interval will probably elapse before the commencement of a new series, and it is at present a subject of consideration whether the Illustrations will be continued in the same form. It has been suggested that monographs of Genera, with the figures on a reduced scale, could be published at a much smaller expense, in proportion to the number of species, and would be more generally useful than the present plan. In compliance with these suggestions, the proprietor is engaged in preparing monographs of several Genera, which will make their appearance at about the fall of the year, and the calculations made give reason to hope that the figures and descriptions may be produced at the rate of about one penny for each species.

From the fact that the figures contained in the present work, are constantly quoted by the English and Continental conchologists, who have written during the course of its publication, there can be no doubt of the desirableness of continuing to publish figures of species, accompanied by information respecting localities, characters, and nomenclature, and the artist would feel a pleasure in doing so, even had he no reason to hope for a greater amount of usefulness or of success than at present enjoyed; but the question is, whether, by altering the plan, and reducing the scale of the figures, he might not command a greater degree of success, and contribute to the advancement of conchological knowledge, to a wider extent. The experiment is about to be made, and in taking leave of the subscribers for a time, the gratifying hope is entertained, that the future re-appearance of the artist, will take place under the influence of their continued encouragement, strengthened by that of many other friends to the science.

TABLE OF CONTENTS.

(The word "*all*" placed against figures in the 3rd column, signifies that the whole of the known species are represented.

The letters "*re*" distinguish those catalogues, which have been re-printed, and of which the old copies are to be cancelled.)

Genera	Numbers of parts				Species and varieties figured	No. of figures	Letter Press	No. of pages
Amphidesma	17	18	19	12	71	Catalogue and Index	2 re.	
Cardium	{ 46 to 51 } 149 150 } 177 to 184 }	70	89	Catalogue and Index	8			
Nucula	14	15	16	24	25	Catalogue and Index	4 re.	
Chiton	{ 38 to 45 }	150	198	{ Catalogue and Index } { Desc. of British species }	10 8			
Chitonellus	{ 159 to 176 }	77	88	Catalogue and Index	10			
Fissurella	68 to 78	80	77	88	Catalogue and Index	10		
Bulinus	{ 21 22 23 26 } 27 30 31 34 } 35 137 to 146 } 185 186 }	119	152	Index, localities, &c	4 re.			
Chilina	135	136	13	all	21all	Catalogue and Index	2 re.	
Neritina	{ 86 87 90 91 } 94 to 100 }	66 all	125all	Catalogue and Index	4 re.			
Eulima	52	53	19 all	19all	Catalogue and Index	2 re.		
Margarita	132	133	134	15 all	30all	Catalogue	2	
Cancellaria	9 to 13	45	47	Catalogue and Index	10 re.			
Murex	{ 58 to 67 } 187 to 199 }	112	128	Catalogue and Index	12			
Typhis	200	6 all	11all	Catalogue and Index	2			
Ranella	{ 84 85 88 } 89 92 93 }	25	37	Index, localities, &c	2			
Monoceros	79	81 to 83	16	27	Catalogue and Index	2 re.		
Eburna	2	20	4	5	Catalogue and Index	2 re.		
Erato	{ 1 to 8 } 101 to 131 }	{ 7 } 4 }	{ 9 } 6 }	{ Catalogue and Index } { to Cyprædæ }	26			
Ovulum								
Cypræa								
Conus	{ 24 25 28 29 } 32 33 36 37 } 54 to 57 } 147 148 } 151 to 153 }	137	153	Index, localities, &c	4			

DIRECTIONS TO THE BINDER.

The lists printed on small paper and published with each part are to be cancelled.

The Plates, Catalogues, Indices, &c. should follow each other in the order indicated in the "Table of Contents."

The word at the head of each plate will point out the genus to which it belongs, and the figures connected with the subjects will shew the order in which the plates of each genus follow each other. They should be placed, so as to lie open on the right side of the book.

The former edition of those Catalogues which have been reprinted, in consequence of having been printed originally on small paper, must be cancelled. The reprints are contained in No. 200,

A CATALOGUE

OF THE

RECENT SPECIES OF THE GENUS


AMPHIDESMA.

1. *variegatum* Lam. Spec. Conch. *Amphidesma*. f. 1, 2, 3.
Brazil.
2. *pulchrum* nob. Z. P. 1832, Dec. Conch. Ill. f. 2. St.
Elena. W. Columbia. var. f. 2* Panama. Mr. Cuming.
3. *pallidum* nob. Z. P. 1832, Dec. Conch. Ill. f. 3. Salango,
W. Col. Mr. Cuming.
4. *formosum* nob. Z. P. 1832, Dec. Conch. Ill. f. 8. St.
Elena. Mr. Cuming.
5. *roseum* nob. Z. P. 1832, Dec. Conch. Ill. f. 1. Tumbes,
Peru. Mr. Cuming.
6. *ellipticum* nob. Z. P. 1832, Dec. Spec. Conch. f. 17.
Monte Christi, W. Col. Mr. Cuming.
7. *australe* nob. Z. P. 1832, Dec. Conch. Ill. f. 4. Lord
Hood's Island, S. Seas, Mr. Cuming, and N. S. Wales,
G. Humphrey.
8. *corrugatum* nob. Z. P. 1832, Dec. Spec. Conch. f. 18.
Iquiqui, Peru. Mr. Cuming.
9. *subtruncatum* nob. Spec. Conch. f. 5, 19, 20. West
Indies.
10. *solidum* Gray, Spicilegium Zoologicum part 1, p. 6, t. 6,
f. 6, Spec. Conch. f. 6. Valparaiso, Mr. Cuming.
11. *cordiforme*, nob. Spec. Conch. f. 7. *Tellina Cordiformis*
Chemn. XI. f. 1931, 1932. Indian Ocean.
12. *Rupium* nob. Z. P. 1832, Dec. Conch. Ill. f. 10. Lord
Hood's Island. var. f. 10* Galapagos Islands. Mr.
Cuming.
13. *cancellatum* nob. Spec. Conch. f. 8. Antigua and St.
Vincent.

14. *lamellosum nob.* Spec. Conch. f. 9, 19. Indian Ocean.
15. *reticulatum nob.* Spec. Conch. f. 11, 12. Brazil and W. Indies.
16. *crenulatum nob.* Spec. Conch. f. 13.
17. *duplicatum nob.* Spec. Conch. f. 14, 15. Ceylon.
18. *punctatum nob.* Z. P. 1832, Dec. Conch. Ill. f. 7. Galapagos Islands. Mr. Cuming.
19. *purpurascens nob.* Z. P. 1832, Dec. Conch. Ill. f. 5. St. Elena. W. Col. Mr. Cuming.
20. *multicostatum nob.* Spec. Conch. f. 16.
21. *lenticulare nob.* Z. P. 1832, Dec. Conch. Ill. f. 9. St. Elena, W. Col. Mr. Cuming.
22. *læve nob.* Z. P. 1832, Dec. Conch. Ill. f. 6 Xipixapi. W. Col. Mr. Cuming.
23. *prismaticum Lam.* Spec. Conch. f. 21. British.
24. *tenue Lam.* Spec. Conch. f. 22. British.
25. *Boysii Lam.* Spec. Conch. f. 23. British.
26. *Siculum nob.* Spec. Conch. f. 24. Sicily.

LIST OF FIGURES. Parts 17 to 19.

- | | |
|------------------------------|-----------------------------|
| Fig. 1. <i>A. roseum.</i> | Fig. 6. <i>A. læve.</i> |
| 2. <i>A. pulchrum.</i> | 7. <i>A. punctatum.</i> |
| 2.* <i>A. pulchrum, var.</i> | 8. <i>A. formosum.</i> |
| 3.* <i>A. pallidum.</i> | 9. <i>A. lenticulare.</i> |
| 4. <i>A. australe.</i> | 10. <i>A. Rupium.</i> |
| 5. <i>A. purpurascens.</i> | 10.* <i>A. Rupium, var.</i> |

Amphidesma.

1. *Cardium* ... C. 1, 12. ...
 2. *Cardium* ... C. 1, 12. ...
 3. *Cardium* ... C. 1, 12. ...
 4. *Cardium* ... C. 1, 12. ...
 5. *Cardium* ... C. 1, 12. ...
 6. *Cardium* ... C. 1, 12. ...



Amphidesma


5




7



6




5



Scapharca



Amphidesma



Amphidromus



CARDIUM.

A

CATALOGUE OF RECENT SPECIES,

BY

By G. B. SOWERBY, Jun.

No.

- 1 *C. costatum*, *Linn. Syst. Nat.* p. 1121. *Chemn. Conch.* t. 15, f. 151. 152. Africa.
- 2 *C. indicum*, *Lam. Anim. sans vert.* t. 6, p. 4. *Con. Illus.* f. 46. N.W. coast of America.
- 3 *C. ringens*, *Lam. Brug. Dict.* No. 21. *Chemn. Conch.* t. 16, f. 170. Gambia, Africa.
- 4 *C. Sinense*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.* f. 35. China.
- 5 *C. Asiaticum*, *Brug. Dict.* No. 19. *Chemn. Conch.* t. 15, f. 153, 154, Nicobar Isl.
- 6 *C. laticostatum*, *Sow. Zool. Proc.* 1833. p. 85. *Con. Illus.* f. 30. Xipixapi, West Colombia. Mr. Cuming.
- 7 *C. fimbriatum*, *Lam. Anim. sans vert.* t. 6. p. 5. *Beech. Zool.* t. xlii. f. 1. *Wood Gen. Conch.* t.—. China.
- 8 *C. tenui-costatum*. *Lam. Anim. sans vert.* t. 6. p. 5. *Con. Illus.* f. 62. (tenue) f. 36. (varium.) f. 19. Swan River, N. Holland.
- 9 *C. striatulum*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.* f. 16. 45. Australia, New Zealand, Mr. G. Benett.
- 10 *C. Australe*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.* f. 12* (*C. Sauciatum*, Beck.—?) Australia, (Var.) China, Ceylon.
- 11 *C. Cumingii*. *Brod. Zool. Proc.* 1833. p. 82. *Con. Illus.* f. 5. Central Amer. G. of Dulce. Mr. Cuming.
- 12 *C. apertum*, *Brug. Dict.* No. 22. *Chem. Conch.* t. 18. f. 181-3. Jamaica.

- No.
- 13 *C. papyraceum*, *Brug.* Dict. No. 29. Chemn. Conch. t. 18, f. 184, Con. Illus. f. 55. 56. Japan.
 - 14 *C. bullatum*, *Lam.* Anim. sans vert. t. 6, p. 6. Solen bullatus, *Linn.* *C. Soleniforme*, *Brug.* Chem. Conch. t. 6. f. 49. 50. Jamaica, St. Vincents.
 - 15 *C. aspersum*, *Sow.* Zool. Proc. 1833. p. 85. Con. Illus. f. 15. St. Elena. Mr. Cuming.
 - 16 *C. ringiculum*. *Sow. Jun.* Zool. Proc. 1840. Con. Illus. f. 11. St. Vincents.
 - 17 *C. latum*, *Brug.* Dict. No. 33. Chemn. t. 19, f. 192. 193. Asia.
 - 18 *C. Pseudolima*, *Lam.* Anim. sans. vert. t. 6, p. 7. Con. Illus. f. 67. Red Sea.
 - 19 *C. Icelandicum*, *Gmel.* p. 3252. Chemn. t. 19, f. 195. 196. Iceland.
 - 20 *C. punctulatum*, *Brod. & Sow.* Zool. Journ. iv. p. 366. Con. Illus. f. 7. Loc. not known.
 - 21 *C. rusticum*, *Linn.* Syst. Nat. p. 1124. Chemn. Conch. t. 19. f. 197. test. Jun. *C. fasciatum*, *Mont.* Test. Brit. Suppl. p. 30, t. 27, f. 6. Britain & Mediterranean.
 - 22 *C. edule*. *Linn.* Syst. Nat. p. 1125, Chemn. Conch. t. 19, f. 194. var. *C. crenulatum*. *Lam.* Britain.
 - 23 *C. nodosum*, Montagu, Test. Brit. p. 81. Turton. Brit. Biv. pl. 13, f. 8. Britain.
 - 24 *C. ovale*, *Nob.* *C. elongatum* (preoccupied) *Mont.* Test. Brit. p. 82. Turt. Brit. Biv. t. 13, f. 9. Britain.
 - 25 *C. exiguum*, *Gmel.* 3255, Lister. t. 317, f. 154. Britain.
 - 26 *C. Siculum*, *Sow. jun.* Zool. Proc. 1840. Con. Illus, f. 31. Sicily.
 - 27 *C. Arcticum*, *Sow. jun.* Zool. Proc. 1840. Con. Illus. f. 26. Arctic Seas.
 - 28 *C. tuberculatum*, *Linn.* Syst. Nat. p. 1122. Chemn. Conch. t. 17, f. 173. Britain and Mediterranean.
 - 29 *C. paucicostatum*, *Sow. jun.* Zool. Proc. 1840. Con. Illus. f. 20. Sicily.

- No.
 30 *C. Deshayesii*, *Payr.* p. 56, t. 1, f. 33. 35. Con. Illus. f. 70. Sicily.
 31 *C. echinatum*, *Linn. Syst. Nat.* p. 1122. Chemn. Conch. t. 200, f. 1951-3. test. jun. *C. ciliare*. Gmel. 29. Britain and Sicily.
 33 *C. aculeatum*. *Linn. Syst. Nat.* p. 1122. Chemn. Conch. t. 15, f. 156. Europe.
 34 *C. erinaceum*. *Lam. Anim. sans vert.* t. 6. p. 8. Chemn. Conch. t. 15. f. 157. Mediterranean.
 35 *C. multispinosum*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 38, 38. a. Misamis, Isl. Mindinao. Mr. Cuming.
 36 *C. exasperatum*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 37. Swan River.
 37 *C. papillosum*, *Poli Test. utriusque Siciliae*, Con. Illus. f. 69. Sicily.
 38 *C. graniferum*, *Brod. and Sow. Zool. Journ.* iv. p. 367. Con. Illus. f. 17. Pacific Ocean. Mr. Cuming.
 39 *C. Isocardia*, *Linn. Syst. Nat.* p. 1122. Chemn. Conch. t. 17, f. 174, 176. America.
 40 *C. Consors*. *Sow. Zool. Proc.* 1833. p. 85. Con. Illus. f. 8. St. Elena. Guacomayo, Mr. Cuming. The imbrications on the ribs are closer than in *C. Isocardia*.
 41 *C. Belcheri* *Brod. & Sow. Zool. Journ.* iv. p. 366. Tab. ix. f. 3. Coast of California and Panuma.
 42 *C. muricatum*, *Linn. Syst. Nat.* p. 1123. Chemn. Conch. t. 17, f. 177. West Indies.
 43 *C. senticosum*, *Sow. Zool. Proc.* 1833. p. 84. Con. Illus. f. 10. St. Elena. Rounder than *C. muricatum*, and tubercles not so oblique. Mr. Cuming.
 43.**C. Radula*, *Brod. & Sow. Zool. Journ.* IV. p. 336. We have not seen this shell. It is in Mr. Bland's collection.
 44 *C. foveolatum*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 65. Swan River.

- No.
- 45 *C. variegatum*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 57. Ribs much closer than in *C. muricatum*.
- 46 *C. unicolor*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 29. 42. Brazils.
- 47 *C. multistriatum*, *Sow. Zool. Proc.* 1833. p. 85. Con. Illus. f. 59. St. Elena, Mr. Cuming.
- 48 *C. maculosum*, *Wood. Gen. Conch.* p. 218, t. 52, f. 3. Con. Illus. f. 63. Ceylon.
- 49 *C. impositum*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 6. 66. China.
- 50 *C. Oxygonum*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 9. China.
- 51 *C. angulatum*, *Lam. Anim. sans vert. Seba. Mus.* 3, t. 86, f. 6. America? The grooves of this common species are deep; the ribs angular and serrated; the shell oval, ventricose.
- 52 *C. marmoreum*, *Lam. Anim. sans vert. Chemn. Conch.* t. 17, f. 179. Jamaica. Flatter than the last, with the anterior ribs smoother, and not so deep.
- 53 *C. Orbita*, *Brod. & Sow. Zool. Proc.* 1833. p. 85. Con. Illus. f. 13. Isl. Annaa, Pacific. O. More oblique, more closely tuberculated and ribbed than *C. marmoreum*. Mr. Cuming.
- 54 *C. elongatum*, *Lam. Brug. Dict.* No. 26. Con. Illus f. 49. (not the figures doubtfully referred to by *Brug. & Lamarck*.) More ventricose, and elongated than the two last, with 39 or 40 flattened ribs; Ticao, Philippines, Mr. Cuming.
- 55 *C. subelongatum*, *Sow. jun. Zool. Proc.* 1840. Con. Illus. f. 61. Ribs fewer and wider apart than in the last.
- 56 *C. maculatum*, (Altered from *maculosum*, preoccupied) *Sow. Zool. Proc.* 1833. p. 85. Con. Illus. f. 18. California. Ribs flattened, nearly smooth; shell not so oblique as *C. subelongatum*.

No.

- 57 *C. enode*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.* f. 51.
Shell more spread, less ventricose, with ribs flatter
and closer than in *C. elongatum*.
- 58 *C. rugosum*, *Lam. Anim. sans vert*, t. 6, p. 10. *C.*
magnum, *Chemn. Conch.* t. 19, f. 191. Yellow
var. *C. flavum*. Linn? *Con. Illus.* (*C. subrugosum*)
f. 32. Dark spotted, *Con. Illus.* f. 39. Pink
spotted, *Con. Illus.* f. 41. Ceylon, Java.
- 59 *C. subrugosum*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.*
f. 34, 71. Dr. Sibbald. Ribs less elevated, shell
less spread, anterior side smoother than the last.
- 60 *C. alternatum*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.*
f. 64. Ticao, Philippines. On the Coral Reefs. Mr.
Cuming.
- 61 *C. procerum*, *Sow. Zool. Proc.* 1833. p. 83. *Con. Illus.*
f. 23. Real Llejos, Central America. Mr. Cuming.
- 62 *C. Panamense*. *Sow. Zool. Proc.* 1833. p. 85. *Con. Illus.*
f. 21. Panama. Mr. Cuming. Ribs fewer and more
angulated than in *C. procerum*.
- 63 *C. ventricosum*, *Brug. Dict.* 25. *C. magnum*, *Born.*
E. M. pl. 299. f. 1. Carolina.
- 64 *C. sulcatum*, *Lam. Anim. sans vert.* tom. 6. p. 10. *C.*
oblongum, *Chemn. Conch.* t. 19. f. 90. *C. flavum*,
Born. Mediterranean. A shell has been collected
at Plymouth by Mr. Cuming, which is distinct
from *C. serratum*, and may probably turn out to
be a British variety of this species.
- 65 *C. elatum*, *Sow. Zool. Proc.* 1833. p. 84. *Con. Illus.*
f. 3. California.
- 66 *C. serratum*, *Linn. Syst. Nat.* 1123. *E. M.* 299. f. 2. *C.*
laevigatum. *Wood. pl.* 54, f. 1. Britain.
- 67 *C. Groenlandicum*, *Gmel. P.* 3252. *Chemn. Conch.* t. 19,
f. 148. var. *edentula*. *Sow. gen. of Sh. Greenland.*
- 68 *C. biradiatum*, *Brug. Dict.* 28. *Chemn. Conch.* t. 18,
f. 185, 186. Ceylon.

- No.
 69 *C. attenuatum*, *Sow. jun. Zool. Proc.* 1840. *lævigatum*,
 var. *biradiatum*, *Wood, Gen. Conch.* pl. 54, f. 2.
 Very much more elongated than the last.
 Ceylon.
- 70 *C. multipunctatum*, *Sow. Zool. Proc.* 1833. p. 84. *Con.*
Illus. f. 14. China.
- 71 *C. lævigatum*, *Linn. Syst. Nat.* p. 1123. *Chemn. Conch.*
 t. 18, f. 189. *C. citrinum*, *Gmel. Wood Gen.*
Conch. pl. 54, f. 3. Atlantic. This name is
 erroneously applied by some authors to the British
C. serratum.
- 72 *C. Brasilianum*, *Lam. Anim. sans vert.* t. 6, p. 5. *Con.*
Illus. f. 60. Brazil.
- 73 *C. Elenensis*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.*
 f. 58. St. Elena. Not so much elongated poste-
 riorly as the last; not lineated. Mr. Cuming.
- 74 *C. Mortoni*, *Conrad. Boston journal?* *Con. Illus.* f. 53,
 54. America. More rounded than *C. Elenensis*,
 and nearly equilateral.
- 75 *C. fragile*, *Sow. jun. Zool. Proc.* 1840. *Con. Illus.* f. 68.
- 76 *C. Æolicum*, *Born. Mus. Cæs.* p. 36. *C. pectinatum*,
Linn? *Chemn. Conch.* t. 18, f. 187, 188. Indian
 Ocean, China.
- 77 *C. lyratum*, *Sow. Zool. Proc.* 1840. *Con. Illus.* f. 40.
 Guinea. This species has no smooth surface on
 the posterior side, as in *C. Æolicum*.
- 78 *C. lineatum*, *Lam. Anim. sans vert.* v. 6, p. 17. *Con.*
Illus. f. 43. America.
- 79 *C. Donaciforme*, *Schroeter, Einl. Conch.* t. 3, p. 68. *Con.*
Illus. f. 27. Asiatic Ocean. Tranquebar.
- 80 *C. parvum*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.*
 f. 33.
- 81 *C. medium*. *Linn. Syst. Nat.* 1122. *Chemn. Conch.* t. 16,
 f. 162, 164. St. Vincents.

- No.
82 *C. biangulatum*, *Sow.* Zool. Journ iv. p. 367. Con. Illus.
f. 2. St. Elena, Mr. Cuming.
83 *planicostatum*, *Sow.* Zool. Proc. 1833. p. 83. Con. Illus.
f. 25. Guacomayo. Mr. Cuming.
84 *C. fornicatum*, *Sow. Jun.* Zool. Proc. 1840. Con. Illus. f. 50.
85 *C. imbricatum*, *Sow. Jun.* Zool. Proc. 1840. Con. Illus.
f. 48. Swan River.
86 *C. Unedo*, *Linn.* Syst. Nat. p. 1123. Chemn. Conch. t. 16,
f. 168, 169. Ceylon.
87 *C. Fragum*, *Linn.* Syst. Nat. p. 1123. Chemn. Conch. t.
16, f. 166. 167. Red Sea.
88 *C. Hemicardium*, *Linn.* Syst. Nat. p. 1121. Chemn.
Conch. t. 16, f. 159, 161. Indian Ocean.
89 *C. tumoriferum*, *Lam.* Anim. sans vert. t. 6, p. 15. Con.
Illus. f. 22. New Holland. Ribs narrower and
tubercles smaller than in the last.
90 *C. obovale*, *Sow.* Zool. Proc. 1833. p. 84. Con. Illus. f. 4.
Xipixapi. Mr. Cuming.
91 *C. subretusum*, *Sow. jun.* Zool. Proc. 1840. p. ¹¹⁰~~84~~. Con.
Illus. f. 24.
92 *C. retusum*, *Linn.* Syst. Nat. p. 1121. Chemn. Conch.
t. 14, f. 139, 140. Suez, Arabia.
93 *C. Auricula*, *Forskäll.* Fauna Arabica. f. 122, No. 52.
Con. Illus. f. 47. Suez, Arabia.
94 *C. Dionæum*, *Sow.* Zool. Journ. iv. p. 367. Beech. Zool.
t. 42, f. 6. 6. Con. Illus. f. 52. S. Pacific Ocean.
95 *C. Cardissa*, *Linn.* Syst. Nat. p. 1121. *Sow.* gen. of Sh.
f. 5. test. jun. (*C. Dionæum*) Con. Illus. f. 1. 1.*
(*C. unimaculatum*,) f. 1.** S. Pacific Ocean. *Nearly*
equilateral.
96 *C. Junoniæ*, *Lam.* Anim. sans vert. t. 6, p. 17. Chemn.
Conch. t. 14, f. 147, 148. *C. Humanum*, *Wood.*
indian Ocean. *Posterior side concave*.
97 *C. inversum*, *Lam.* Anim. sans vert. t. 6, p. 17. Chemn.
Conch. t. 14, f. 149, 150. Indian Ocean. *Anterior*
side concave.

CORRECTED LIST OF FIGURES.

CARDIA ;—Parts 48 to 51, 149—150, 177 to 186.

FIG.		FIG.	
1 1.* 1.**	<i>C. Cardissa</i> , Sp. 95, young.	35	<i>C. Sinense</i> , Sp. 4.
2	<i>C. biangulatum</i> , Sp. 82.	36	<i>C. tenuicostatum</i> , Sp. 8.
3	<i>C. elatum</i> , Sp. 65.	37	<i>C. exasperatum</i> , Sp. 36.
4	<i>C. obovale</i> , Sp. 90.	38	<i>C. multispinosum</i> , Sp. 35.
5	<i>C. Cumingii</i> Sp. 11.	39	<i>C. rugosum</i> , var. Sp. 58.
6	<i>C. impositum</i> , Sp. 49.	40	<i>C. lyratum</i> , Sp. 77.
7	<i>C. punctulatum</i> , Sp. 20.	41	<i>C. rugosum</i> , var. Sp. 58.
8	<i>C. Consors</i> , Sp. 40.	42	<i>C. unicolor</i> , Sp. 46.
9	<i>C. Oxygonum</i> , Sp. 50.	43	<i>C. lineatum</i> , Sp. 78.
10	<i>C. senticosum</i> , Sp. 43.	44	<i>C. lineatum</i> , young, Sp. 78.
11	<i>C. ringiculum</i> , Sp. 16.	45	<i>C. striatulum</i> , Sp. 9.
12*	<i>C. Australe</i> , Sp. 10.	46	<i>C. indicum</i> , Sp. 2.
13	<i>C. Orbita</i> , Sp. 53.	47	<i>C. Auricula</i> , Sp. 93.
14	<i>C. multipunctatum</i> , Sp. 70.	48	<i>C. imbricatum</i> , Sp. 85.
15	<i>C. aspersum</i> , Sp. 15.	49	<i>C. elongatum</i> , Sp. 54.
16	<i>C. striatulum</i> , Sp. 9.	50	<i>C. fornicatum</i> , Sp. 84.
17	<i>C. graniferum</i> , Sp. 38.	51	<i>C. enode</i> , Sp. 57.
18	<i>C. maculatum</i> , Sp. 56.	52	<i>C. Dionæum</i> , Sp. 94.
19	<i>C. tenuicostatum</i> , Sp. 8.	53, 54	<i>C. Mortoni</i> , Sp. 74.
20	<i>C. paucicostatum</i> , Sp. 29.	55, 56	<i>C. papyraceum</i> , Sp. 13.
21	<i>C. Panamense</i> , Sp. 62.	57	<i>C. variegatum</i> , Sp. 45.
22	<i>C. tumoriferum</i> , Sp. 89.	58	<i>C. Elenense</i> , Sp. 73.
23	<i>C. procerum</i> , Sp. 61.]	59	<i>C. multistriatum</i> , Sp. 47.
24	<i>C. subretusum</i> , Sp. 91.	60	<i>C. Brasilianum</i> , Sp. 72.
25	<i>C. planicostatum</i> , Sp. 83.	61	<i>C. subelongatum</i> , Sp. 55.
26	<i>C. arcticum</i> , Sp. 27.	62	<i>C. tenuicostatum</i> , Sp. 8.
27	<i>C. Donaciforme</i> , Sp. 79.	63	<i>C. maculosum</i> , Sp. 48.
28	<i>C. echinatum</i> , var. Sp. 31.	64	<i>C. alternatum</i> , Sp. 60.
29	<i>C. unicolor</i> , Sp. 46.	65	<i>C. foveolatum</i> , Sp. 44.
30	<i>C. laticostatum</i> , Sp. 6.	66	<i>C. impositum</i> , Sp. 49.
31	<i>C. Siculum</i> , Sp. 26.	67	<i>C. Pseudolima</i> , Sp. 18.
32	<i>C. rugosum</i> , var. Sp. 58.	68	<i>C. fragile</i> , Sp. 75.
33	<i>C. parvum</i> , Sp. 80.	69	<i>C. papillosum</i> , Sp. 37.
34	<i>C. subrugosum</i> , Sp. 59.	70	<i>C. Deshayesii</i> , Sp. 30.
		71	<i>C. subrugosum</i> , Sp. 59.

Cardium.



2



1



2



3




1*



1**



4



Cardium.

5



6



7



8



9



10

EXHIBITION



Cardium.


12



11



12*




13



14



15



Faint, illegible handwritten text or markings.

Cardium.



16



17



18



19




20




21



Cardium.



22



22



23



24



24



25



11



12



13




14




15

Cardium.


26



27



27



28



29




31



31



30



Cardium.


32



33



35



34



36

Cardium




Cardium.

37




37



38.a




38.b



39

Cardium



Cardium



14



14



10



15



15



16



Cardium


43



41






42




40

Conchus




Cardium


Cardium



Cardium



Cardium




Cardium



58



58



55



57



56

1846



Cardium

60



59



61



63



62

Cardium



22



23



24



25



26

Cardium


64



65




66



67

Cardium



107.
Cardium



69




68



62



70




70



71

Curculion



A CATALOGUE OF RECENT SPECIES OF NUCULÆ

* *LANCEOLATÆ*.

- 1 *lanceolata* *Lam.* Sowerby Gen. of Shells, No. 17, f. 1.
Conch. Illustr. f. 1. Xipixapi.
- 2 *Elongata nobis.* Z. P. 1832, Dec. Conch. Illustr. f. 2.
Salango, Mr. Cuming.
- 3 *Tellinoides nobis*, Gen. of Shells, No. 17, f. 2. Cumana.
- 4 *Crenifera nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 3.
Xipixapi, Mr. Cuming.

** *ROSTRATÆ*.

- 5 *arctica*, *Brod. and Sow.* IV. pl. IX. f. 1. Kamtschatka.
- 6 *Nicobarica* *Lam.* Conch. Illustr. f. 4.
- 7 *curvirostrum*, Humphrey M. S. Conch. Illustr. f. 5.
Shell oblong, thin, pellucid, white, smooth, rather gibbous; anterior side rounded, posterior acuminate, turned upward, with three crenated ridges; surface covered with oblique, nearly longitudinal grooves: length 0·7, breadth 0·25, height 0·4. Coast of Guinea, *Humphrey. Arca rostrata* Mont. Test. Brit. Sup. p. 55. t. 27.
- 8 *Pella* *Lam.* Sowerby's Gen. of Shells No. 17. f. 4.
Conch. Illustr. f. 6.
- 9 *glacialis* *Wood* Sup. pl 2. f. 6.
- 10 *fluviatilis nobis*, Sow. Gen. of Shells, No. 17, f. 3.
(*N. rostrata* *Lam.*) Rivers of Coromandel, Schroëtter.
- 11 *minuta* *Mont.* Test. Brit. p. 144. Sowerby Gen. of Shells, No. 17, f. 5. (*N. rostrata* Sowerby) British.
- 12 *parva nobis*, Conch Illustr. f. 7. Shell oblong, white, thickish, with an olivaceous epidermis, longitudinally

grooved; anterior side rounded, posterior rather beaked, angular: dorsal posterior area not grooved; length 0.5, breadth 0.2, height 0.3. The only specimen I have seen was brought from Melville Island in 1824, by Captain Parry.

- 13 *costellata nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 8. Panama. Mr. Cuming.
- 14 *gibbosa nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 9. Gulf of Nocoiyo. Mr. Cuming.
- 15 *eburnea nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 10. Bay of Caraccas. Mr. Cuming,
- 16 *polita nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 11. Panama. Mr. Cuming.
- 17 *nasuta, nobis*, Conch. Illustr. f. 12. Shell oblong, white, very finely longitudinally striated; anterior side rather shorter than the posterior, rounded; posterior acuminate, slightly turned upwards, with an elongated longitudinally striated dorsal area: length 0.8, breadth 0.33, height 0.4. From the East Indies. G. Humphrey.
- 18 *Fabula, nobis*, Conch. Illustr. f. 13. Shell ovate, white, obliquely striated, nearly equilateral, anterior side rounded, posterior acuminate, slightly turned upwards, with two ridges passing from the umbones to the acuminate extremity, the upper of which is crenated, forming a dorsal posterior area: length 0.4, breadth 0.15, height 0.25.
- 19 *Elenensis nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 14. St. Elena, Mr. Cuming.
- 20 *cuneata nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 15. Valparaiso. Mr. Cuming.
- 21 *striata nobis*, Conch. Illustr. f. 16. Shell ovate, white, concentrically striated, nearly equilateral, anterior and posterior dorsal slopes nearly equal; anterior side slightly acuminate, rounded; posterior side acuminate, with an indistinct dorsal area: length 0.55, breadth 0.3, height 0.4. From China, G. Humphrey.

22 *Mauritiana nobis*, Conch. Illustr. f. 17. Shell ovate white, concentrically grooved, nearly equilateral, anteriorly rounded, posteriorly acuminate; dorsal slope nearly equal on both sides; length 0.4, breadth 0.2 height 0.25. From the Mauritius, G. Humphrey.

23 *limatula* Say, American Conchology t. 12.

24 *lævis* Say, Amer. Conch. t. 12.

25 *concentrica* Say, Amer. Conch. t. 12.

*** *ELLIPTICÆ*.

26 *margaritacea* Lam. Sow. Gen. of Shells, f. 7. British.

27 *decussata nobis*, Conch. Illustr. f. 18. Shell somewhat triangular, white, decussately striated; anterior side shorter, its slope rather elevated in the middle; posterior side much longer, oblique; ventral margin rounded its edge crenated; length 0.85, breadth 0.4, height 0.6. Brought from the Coast of Guinea, in 1788, G. Humphrey.

28 *rugulosa nobis*, Conch. Illustr. f. 19. Shell ovate triangular, with a greenish epidermis, covered with rugulose striæ; angles obtuse; anterior slope with a central elevation; posterior slope somewhat rounded; ventral margin very finely crenulated; length 0.5, breadth 0.25, height 0.4.

29 *nitida nobis*, Conch. Illustr. f. 20. Shell ovate sub-triangular, smooth, shining, white, with an olive coloured epidermis; posterior slope somewhat rounded, anterior nearly straight, with a central elevation; length 0.5, breadth 0.25, height 0.4. Common on the British Coast.

30 *obliqua* Lam. Conch. Illustr. f. 21. Valparaiso, Mr. Cuming.

31 *tenuis* Mont. Test. Brit. Sup. p. 56. t. 29. f. 1. Scarborough, Frith of Forth, &c.

32 *convexa nobis*, Conch. Illustr. f. 22. Shell ovate, turgid, whitish, smooth; anterior side shorter, slightly acumi-

nated ; posterior side much longer, dorsal slope rounded ; ventral margin entire ; hinge pit very oblique, teeth of anterior side very few ; length 0.55, breadth 0.35, height 0.4. From China, G. Humphrey.

33 *Pisum. nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 23, Valparaiso. Mr. Cuming.

34 *exigua nobis*, Z. P. 1832, Dec. Conch. Illustr. f. 24. and magnified f. 24*. Bay of Caraccas. Mr. Cuming.

LIST OF FIGURES.

- | | |
|-------------------------------|-----------------------------|
| Fig. 1. <i>N. Lanceolata.</i> | Fig. 13. <i>N. Fabula.</i> |
| 2. <i>N. elongata.</i> | 14. <i>N. Elenensis.</i> |
| 3. <i>N. crenifera.</i> | 15. <i>N. cuneata.</i> |
| 4. <i>N. nicobarica.</i> | 16. <i>N. striata.</i> |
| 5. <i>N. curvirostra,</i> | 17. <i>N. Mauritiana.</i> |
| 6. <i>N. Pella.</i> | 18. <i>N. decussata.</i> |
| 7. <i>N. parva.</i> | 19. <i>N. rugulosa.</i> |
| 8. <i>N. costellata.</i> | 20. <i>N. nitida.</i> |
| 9. <i>N. gibbosa.</i> | 21. <i>N. obliqua.</i> |
| 10. <i>N. eburnea.</i> | 22. <i>N. convexa.</i> |
| 11. <i>N. polita.</i> | 23. <i>N. Pisum.</i> |
| 12. <i>N. nasuta.</i> | 24. <i>N. exigua.</i> |
| | 24.* <i>N. exigua. mag.</i> |

Nucula.

1



2



3



Nucula.

Alvada



Nucula

22



20



19



18



23



24x



24



21



A CATALOGUE

OF THE

RECENT SPECIES OF CHITONES.

No.

- 1 Ch. *Echinatus*, *Barnes*, Amer. Journ. of Sc. vol. 7, No. 1, 1823. *Tuberculiferus*. *Sow.* Tanc. Cat. *Spiniferus*, *Frembly*, Zool. Journ. vol. 3, p. 196, Suppl. pl. xvi. f. 1. *Spiniferus*, Var. Con. Illus. f. 47, Peru.
- 2 Ch. *Spinosus*, *Brug.* Journ. D'Hist. Nat. vol. 1, p. 25. *Sow.* Gen. Chiton. f. 1, Con. Illus. f. 151, Australia.
- 3 Ch. *Spiniger*, *Nob.* Mag. Nat. Hist.* 1840, Con. Illus. f. 68.
- 4 Ch. *Brevispinosus*, *Nob.* Mag. Nat. Hist. 1840, Con. Illus. f. 136. Island Johanna.
- 5 Ch. *Blainvillii*, *Brad.* Zool. Proc. 1832, p. 27. Con. Illus. f. 6. Inner Lobos Island.
- 6 Ch. *Rugosus*, *Gray* ———? Con. Illus. f. 49.
- 7 Ch. *Alternatus*, *Nob.* Mag. Nat. Hist. Con. Illus. f. 141. Japan.
- 8 Ch. *Pectinatus*, *Nob.* Mag. Nat. Hist. Con. Illus. f. 146.
- 9 Ch. *Loo-chooanus*, *Sow.* Zool. Journ. vol. iv, p. 368, Loo-choo.
- 10 Ch. *Piceus*, *Gmel.* Linn, p. 3204, Wood, Gen. Con. t. 1, f. 3. *convexus*, *granulatus*, *gemmaus*, tu-

* Charlesworth's New Series of Lowdon's Magazine of Natural History for June, 1840.

No.

- berculatus, unguiculatus, zonatus, *Auct.* Con. Illus. f. 147.
- 11 Ch. Magnificus, Gray? *Con. Illus.* f. 52.
- 12 Ch. Niger, *Barnes*, Amer. Journ. Sc. vol. 7, No. 1, 1833, p. 671. Coquimbensis, *Frembl.* Zool. Journ. vol. iii, p. 197. Tab. Suppl. xvi, f. 2. Peru.
- 13 Ch. Squamosus, *Linn.* Syst. Nat. 1107, Wood, Gen. Con. pl. 1, f. 1, Var. *Bistriatus*, Wood, Chemn. Conch. 8, t. 94, f. 788, 791, Var. Obscurus. *Nob.* Con. Illus. f. 76. West Indies.
- 14 Ch. Sículus, *Gray*, Spic. Zool. 1, p. 5. Con. Illus. f. 123 to 125. *Pollii* Deshayes. Sicily.
- 15 Ch. Lyratus, *Nob.* Mag. Nat. Hist. Con. Illus. f. 126.
- 16 Ch. Foveolatus, *Nob.* Mag. Nat. Hist. Con. Illus. f. 60.
- 17 Ch. Excavatus, Gray? Con. Illus. f. 131.
- 18 Ch. Fasciatus, Wood, Gen. Con. p. 10, Con. Illus. f. 153. South America.
- 19 Ch. Australis, *Nob.* Mag. Nat. Hist. Con. Illus. f. 46. Australia.
- 20 Ch. Stokesii, *Brod.* Zool. Proc. 1832, p. 25, Con. Illus. f. 24. Columbia.
- 21 Ch. Sulcatus, Wood, Gen. Con. p. 16, Con. Illus. f. 12.
- 22 Ch. Virgulatus, *Nob.* Mag. Nat. Hist. Con. Illus. f. 132.
- 23 Ch. Patulus, *Nob.* Mag. Nat. Hist. Con. Illus. f. 134.
- 24 Ch. Granulosus, *Frembl.* Zool. Journ. vol. iii, p. 201. Tab. Suppl. xvii, f. 3. Conception Bay.
- 25 Ch. Granosus, *Frembl.* Zool. Journ. vol. iii, p. 200. Tab. Suppl. xvii, f. 1. Chili.
- 26 Ch. Barnesii, Gray, Spic. Zool. 1, p. 5, Con. Illus. f. 2. Coquimbo.

- No.
- 27 Ch. *Marmoratus*, *Gmel.* p. 3205, *Con. Illus.* f. 148, 149, 150. West Indies.
- 28 Ch. *Cymbiola*, *Nob. Mag. Nat. Hist. Con. Illus.* f. 45, 85, 86, *Tulippa*, Quoy, C. of G. H.
- 29 Ch. *Articulatus*, *Sow. Zool. Proc.* 1832, 59, *Con. Illus.* f. 18. California.
- 30 Ch. *Lævigatus*, *Sow. Zool. Proc.* 1832, 59, *Con. Illus.* f. 18,* California.
- 31 Ch. *Godallii*, *Con. Illus.* f. 34, 50, *Zool. Proc.* 1832, p. 25. Gallapagos.
- 32 Ch. *Bowenii*, *King, Zool. Journ.* vol. 5, p. 338, *Con. Illus. Chiton*, f. 37, *Terradel Fuego, Str. Magellan*
- 33 Ch. *Quoyi*, *Bl. Dict. des Sc. Oseabrion*, *Ch. viridis*. Quoy et Gaimard, *Voyage de l'Astrolabe*, t. 3, p. 381, pl. 74, f. 17—22.
- 34 Ch. *Striatus*, *Barnes, Amer. Journ. Sc.* vol. 7, No. 1, 1823, *Latus*, *Sow. Tanc. Cat. Olivaceus*, *Frembl. Zool. Journ.* III, p. 199, *Tab. Suppl.* xvi. f. 4, *Chili*, *Var. subfuscus*, *Sow. Zool. Proc.* 1832, p. 26, *Con. Illus.* f. 3, 41, *Chiloe*.
- 35 Ch. *Tenuistriatus* *Nob. Mag. Nat. Hist. Con. Illus.* f. 135,
36. Ch. *Lentiginosus*, *Nob. Mag. Nat. Hist. Con. Illus.* 120—1.
- 37 Ch. *Concinnus*, *Nob. Mag. Nat. Hist. Con. Illus.* 1178. *Chonos*.
- 38 Ch. *Evanidus*, *Nob. Mag. Nat. Hist. Con. Illus.* f. 139.
- 39 Ch. *Georgianus*, Quoy et Gaimard, *Voy. Astrolabe*, t. 3, p. 379, pl. 25 to 30. K. George's Port.
- 40 Ch. *Cumingii*, *Frembl. Zool. Journ.* vol. iii, p. 198. *Tab. Suppl.* xvi, f. 3. *Var. Con. Illus.* 32, 51. *Valparaiso*.

No.

- 41 Ch. *Glauco-cinctus*, Frembl. Zool. Journ. vol. iii, p. 201, Tab. Suppl. xvii, f. 2. Valparaiso.
- 42 Ch. *Albo-lineatus*, Sow. Zool. Journ. vol. 4, p. 368. Con. Illus. f. 39. Mexico.
- 43 Ch. *Versicolor*, Nob. Mag. Nat. Hist. Con. Illus. f. 75, var. 122.
- 44 Ch. *Sculptus*, Nob. Mag. Nat. Hist. Con. Illus. f. 66.
- 45 Ch. *Longicymba*, Bl. Dict. des Sc. *Oscabrion*, Ch. var. Con. Illus. f. 67. Hunter's River.
- 46 Ch. *Textilis*, Gray, Spic. Zool. 1, p. 5, t. 6, f. 20. Ch. *Indicus*, Con. Illus. f. 55, var. Ch. *Solea*, Con. Illus. f. 61, Cape of G. H.
- 47 Ch. *Dispar*, Sow. Zool. Proc. 1832, p. 28, Con. Illus. f. 25, Panama.
- 48 Ch. *Punctulatissimus*, Sow. Zool. Proc. 1832, p. 28. Con. Illus. f. 9, 26. Beechy's Appendix, p. 40, f. 2, 3. Mexillones, Iquiqui. Africa.
- 49 Ch. *Stramineus*. Sow. Zool. Proc. 1832, p. 105. Con. Illus. f. 28, var. 838. Chiloe.
- 50 Ch. *Fimbriatus*, Nob. Mag. Nat. Hist. Con. Illus. f. 137, Peru.
- 51 Ch. *Apiculatus*, Say? Con. Illus. f. 140. Massachusetts Bay.
- 52 Ch. *Lævis*. See Catalogue of British Chitones.
- 53 Ch. *Cancellatus*. Do.
- 54 Ch. *Albus*. Do.
- 55 Ch. *Marginatus*. Do.
- 56 Ch. *Cinereus*. Do.
- 57 Ch. *Latus*. Do. *Fulminatus*, Couthoy, Boston. Amer. Journ.
- 58 Ch. *Limaciformis*, Sow. Zool. Proc. 1832, p. 58. Con. Illus. f. 38. Columbia.

- No.
 59 Ch. *Roseus*, Sow. Zool. Proc. 1832, p. 58. Con. Illus. f. 14. Columbia.
 60 Ch. *Lyellii*, Sow. Zool. Proc. 1832, p. 26. Con. Illus. f. 7. Pitcairn's Island.
 61 Ch. *Catenulatus*, Sow. Zool. Proc. 1832, p. 104. Con. Illus. f. 145. Inner Lobos Island.
 62 Ch. *Rugulatus*, Sow. Zool. Proc. 1832, p. 58, var. in conspicuus, Nob. Con. Illus. f. 42, 143, 144. Inner Lobos Island.
 63 Ch. *Grayi*, Sow. Zool. Proc. 1832, p. 57. Con. Illus. f. 8, 16. Peru.
 64 Ch. *Crenulatus*, Brod. Zool. Proc. 1832, p. 27. Con. Illus. f. 43.
 65 Ch. *Platymerus*, Sow. Zool. Beech, Appendix, p. 149, pl. 41, f. 11.
 66 Ch. *Bipunctatus*, Sow. Zool. Proc. 1832, p. 104. Con. Illus. f. 27, 155.
 67 Ch. *Pusillus*, Sow. Zool. Proc. 1832, p. 57. Con. Illus. f. 31. Peru.
 68 Ch. *Exiguus*, Sow. Zool. Proc. 1832, p. 104. Con. Illus. f. 36. Lord Hood's Island.
 69 Ch. *Fulvus*, Wood, Gen. Con. p. 7, t. 1, f. 2. Con. Illus. f. 83, var. ? (*velatus*,) Con. Illus. f. 53. Cadiz.
 70 Ch. *Swainsoni*, Sow. Zool. Proc. 1832, p. 27. Con. Illus. f. 5. Peru.
 71 Ch. *Lineatus*, Wood, Gen. Con. p. 15, pl. 2, f. 45. Con. Illus. f. 77.
 72 Ch. *Atratus*, Nob. Mag. Nat. Hist. Con. Illus. f. 57, 58. Falkland Island.
 73 Ch. *Castaneus*, Wood, Gen. Con. p. 13, pl. 2, f. 2, 3; pl. 3, f. 2, 3. Con. Illus. f. 114, 115, 116. C. of Good Hope.

No.

- 74 Ch. Gigas, Gmel. p. 3206, Chem. Con. 8, t. 96, f. 819. C. of Good Hope.
- 75 Ch. Elegans, Fremby, *Tab. Suppl.* xvii, f. 6. Con. Illus. 119, f. 13, Valparaiso, *var. Chilensis*, *Fremby*, Zool. Journ. vol. iii, p. 204. *Tab. Suppl.* xvii, f. 8. Chili. *var. Ch. Chiloensis*, *Sow.* Zool. Proc. 1832, p. 58. Con. Illus. f. 13,* 10, 29, 30. Chiloe. *var. Ch. Lineolatus*, f. 154, *Fremby*, Zool. Journ. vol. iii. p. 204. *Tab. Suppl.* xvii, f. 7. Con. Illus. (*var.*) f. 50, 30, Valparaiso, *var. Ch. Graniferus*, *Sow.* Zool. Proc. 1832, p. 104. [Conception Bay, *Young*, *Sparsus*, Con. Illus. f. 73, 74.
- 76 Ch. Fastigiatus, *Gray* ———? Con. Illus. f. 11, 35. California.
- 77 Ch. Disjunctus, *Fremby*, Zool. Journ. vol. iii, p. 203. *Tab. Suppl.* xvii, f. 5. Valparaiso.
- 78 Ch. Janierensis, *Gray* Sp. Zool. p. 6, f. 8. Con. Illus. f. 56, *var. (Occidentalis)*. Con. Illus. f. 54. Rio Janiero.
- 79 Ch. Ellinensis, *Sow.* Zool. Proc, 1832, p. 27. Con. Illus. f. 69. St. Helena.
- 80 Ch. Cajetanus, *Polli*. Test. (Ch. concentricus). Con. Illus. f. 72.
- 81 Ch. Tunicatus, *Wood*, Gen. Con. p. 11, pl. 2, f. 1. Con. Illus. f. 152. Arctic Ocean.
- 82 Ch. Monticularis, *Quoy*, Voy. Astro. t. 3, p. 406, pl. 73, 30. to 36. Con. Illus. f. 129.
- 83 Ch. Vestitus, *Sow.* Zool. Journ. vol. 4, p. 368. Arctic Ocean. Ch. Emersoni, *Couthoy*, Mass. Bay. Con. Illus. f. 128.
- 84 Ch. Amiculatus, *Pallas*, Nov. Act. Petrop, ii, p. 241, t. 7, f. 26—30. Con. Illus. f. 80. Kurile Islands.

- No.
- 85 Ch. (Chitonellus) Striatus, *Lam. Hist. Nat. Anim. sans vert.* vol. 6, p. 317. Con. Illus. f. 62. New Holland.
- 86 Ch. (Chitonellus) Lœvis, *Lam. Anim. sans vert.* vol. 6, p. 517. Sow. Gen. of Sh. New Holland.
- 87 Ch. (Chitonellus) Strigatus, *Nob. Mag. Nat. Hist.* Con. Illus. f. 63.
- 88 Ch. Hastatus, *Nab. Mag. Nat. Hist.* Con. Illus. f. 127.
- 89 Ch. Crinitus. See Cat. of British Chitones.
- 90 Ch. Fascicularis. Do.
- 91 Ch. Hirudiniformis, Sow. Zool. Proc. 1832, p. 59, Con. Illus. f. 23, 48, 142. Garnoti, Bl. Peru. Ch. Danielli. Con. Illus. f. 48, which comes from the Cape of Good Hope, may prove to be a distinct species—but all our Specimens are much worn.
- 92 Ch. Violaceus, *Quoy et Gaimard, Voy. del' Astrolabe*, t. 3, p. 403, pl. 73, f. 15, 16, var. 17, 20, Con. Illus. f. 133.
- 93 Ch. Retusus, Sow. Zool. Proc. 1832, p. 28, Con. Illus. f. 22. Guacomayo.
- 94 Ch. Peruvianus, *Lam. Anim. sans. vert.* Enc. Met. pl. 163, f. 7, 8, var. Zool. Journ. Tab. Suppl. f. 4, var. Con. Illus. f. 44, 44,* 44.** Peru.
- 95 Ch. Setiger, *King, Zool. Journ.* vol. 5, p. 338. Con. Illus. f. 17, var. Ch. Fremblii, *Brod. Zool. Proc.* 1832, p. 28, Con. Illus. f. 4, var. Ch. setosus, Zool. Beechey, Terra del Fuego, C. Biramosus, Quoy, Costatus, Bl.
- 96 Ch. Setosus, Sow. Zool. Proc. 1832, p. 27, Con. Illus. f. 19. Guacomayo.
- 97 Ch. Watsoni, *Nob. Mag. Nat. Hist.* Con. Illus. f. 81, 82, 130. Ch. Castanus, Quoy et Gaimard, Voy.

No.

- Astr. t. 3, p. 387, pl. 74, f. 33—34. C. of Good Hope.
- 98 Ch. Hennahi, *Gray*, Sp. Zool. p. 5, Con. Illus. f. 1, 33. Callao Bay.
- 99 Ch. Spinulosus, *Gray*, Sp. Zool. p. 5, t. 6, f. 7, 7, a. Con. Illus. 70, 71, Rio Janiero. Ch. Columbiensis. Sow. Zool. Proc. Con. Illus. f. 15, Ch. Luridus. Sow. Zool. Proc. Con. Illus. f. 20? St. Helena.
- 100 Ch. Scabriculus, *Sow.* Zool. Proc. 1832, p. 28. Con. Illus. f. 21. Guacomayo.
- 101 Ch. Ciliatus, *Nob.* Mag. Nat. Hist. Con. Illus. f. 79.
- 102 Ch. Petholatus, *Nob.* Mag. Nat. Hist. Con. Illus. f. 64, 65. var. Porphyrius, f. 59. Australia.

N.B.—The remaining Plates, with the Index to the Chitones, will be published for June.

CORRECTED LIST OF FIGURES

In Parts 38 to 45, and 159 to 176.

The Nos. after the Names indicate the situation of the Species in the Catalogue.

Fig.		Fig.	
1	Hennahi, var. 93.	31	Pusillus, 67.
2	Barnesii, 26.	32	Cumingii, 40.
3	Striatus, var. 34.	33	Hennahi, var. 98.
4	Setiger, var. 95.	34	Goodallii, var. 31.
5	Swainsoni, 70.	35	Fastigiatus, 76.
6	Blainvillii, 5.	36	Exiguus, 68.
7	Lyelli, 60.	37	Bowenii, 32.
8	Grayi, 63.	38	Limaciformis, 58.
9	Punctulatissimus, 48.	39	Albolineatus, 42.
10	Elegans, var. 75.	40	Goodallii, 31.
11	Fastigiatus, 76.	41	Striatus, var. 34.
12	Sulcatus, 21.	42	Rugulatus, 62.
13	Elegans, 75.	43	Crenulatus, 64.
13*	——, var. 75.	44	Peruvianus, var. 94.
14	Roseus, 59.	45	Cymbiola, 28.
15	Columbiensis, 99.	46	Australis, 19.
16	Grayi, var. 63.	47	Echinatus, var. sp. 1.
17	Setiger, 95.	48	Danielli, 91.
18	Articulatus, 29.	49	Rugosus, 6.
18*	Lævigatus, 30.	50	Elegans, var. 75.
19	Setosus, 96.	51	Cumingii, var. 40.
20	Luridus, 99.	52	Magnificus, 11.
21	Scabriculus, 100.	53	Fulvus, var. ? 69.
22	Retusus, 93.	54	Janierensis, 78.
23	Hirudiniformis, 91.	55	Textilis ? 46.
24	Stokesii, 20.	56	Janierensis, 78.
25	Dispar, 47.	57, 58.	Atratus, 72.
26	Punctulatissimus, 48.	59	Petholatus, var. 102.
27	Bipunctatus, 66.	60	Foveolatus, 16.
28	Stramineus, 49.	61	Textilis, 46.
29	Elegans, var. 75.	62	(Chitonellus) striatus,
30	—— 75.		85.

Fig.	Fig.	
63	(Chitonellus) strigatus, 126	Lyratus, 15.
	87.	127 Hastatus, 88.
64, 65.	Petholatus, 102.	128 Vestitus, 83.
66	Sculptus, 44.	129 Monticularis, 82.
67	Longicymba, var. 45.	130 Watsoni, var. 97.
68	Spiniger, 3.	131 Excavatus, 17.
69	Elenensis, 79.	132 Virgulatus, 22.
70, 71.	Spinulosus, 99.	133 Violaceus, 92.
72	Cajetanus, 80.	134 Patulus, 23.
73, 74.	Elegans, (Young) 75.	135 Tenuistriatus, 35.
75	Versicolor, 43.	136 Brevispinosus, 4.
76	Squamosus, var. 13.	137 Fimbriatus, 50.
77	Lineatus, 71.	138 Stramineus, var. 49.
78	Rugulatus? 62.	139 Evanidus, 38.
79	Ciliatus, 101.	140 Apiculatus, 51.
80	Amiculatus, 84.	141 Alternatus, 7.
81	Watsoni, 97.	142 Hirudiniformis,
82	————— var. 97.	143 Rugulatus, var. 62.
83	Fulvus, 69.	144 ————— var. 62.
84	Spinulosus, 99.	145 Catenulatus, 61.
85, 86.	Cymbiola, var. 28.	146 Pectinatus, 8.
87 to 113.	British species,	147 Piceus, 10.
	p. 7.	148 Marmoratus, 27.
114	Castaneus, 73.	149 ————— var.
115, 116.	Castaneus, var. 73.	150 ————— var.
117, 118.	Concinnus, 37.	151 Spinosus, sp. 2.
119	Elegans, 75.	152 Tunicatus, 81.
120	Lentiginosus, 36.	153 Fasciatus, 18.
121	————— worn, 36	154 Chilensis, var. Lineatus,
122	Versicolor, var. 43.	75.
123 to 125.	Siculus, 14.	155 Bipunctatus, 66.

DESCRIPTIVE CATALOGUE
OF
BRITISH CHITONES,

By G. B. SOWERBY, JUN.

While engaged in preparing a Catalogue of the genus Chiton for publication, with these illustrations, my attention has been directed to the confusion, prevailing among authors and collectors, respecting the British species. It having consequently occurred to me that effectual service would be rendered to science by determining the nomenclature and synonymes of this section, I have procured specimens from the principal cabinets, through the kindness of their possessors, and compared them carefully with the published figures and descriptions. The following result is presented in a separate form, for the convenience of those collectors whose attention is confined to species found on our own coasts.

Sp. 1. Ch. FASCICULARIS.

Descr. Shell elongated, depressed; valves sub-triangular, coarsely granulated at the sides; with

a dorsal striated band; margin rough, hairy, with bunches of bristles on both sides; inserted laminæ of the valves narrow; colour brown, slightly varied, with one black, and two white, dorsal bands.

C. fascicularis, Linn. Syst. Nat. p. 1106.

Chemn. Conch. 10. t. 173. f. 1688.

Sow. Jun. Conch. Illustr. f. 87. 87 a.

Obs. This species is much more depressed than the following; the valves are closer at the sides, and the granulations coarser. As Linnæus appears not to have recognized two fasciculated species, it cannot be known which he intended to describe; but there can be no doubt of the present species being identical with that figured in Chemnitz.

Sp. 2. *C. CRINITUS*.

Descr. Shell oblong; strongly keeled by a dorsal striated rib; valves beaked, reniform, finely granulated at the sides, disunited at the extremities; margin smooth, hairy, fasciculated; inserted laminæ broad; colour various.

Ch. crinitus, Pennant, Brit. Zool.

Ch. discrepans, Brown.

Ch. fascicularis, Sow. Gen. of Sh. Wood's Gen. Conch. pl. 2. f. 6.

Ch. crinitus, Sow. Jun. Conch. Illustr. f. 88, 93.

Obs. This being the most common of the two fasciculated species, has been most frequently figured under the original name; which, however, it seems to me

most correct to retain for the shell of Chemnitz, his figure being the oldest, and universally quoted. The general appearance and the strong central band of striae in Pennant's figure, leave no doubt on my mind that he intended to represent the shell to which we have fixed the name proposed by him.

Sp. 3. *C. MARGINATUS*.

Descr. Shell oval, carinated. elevated, posteriorly widened, minutely granulated; margin also minutely granulated; colour various.

Ch. marginatus, Pennant, Brit. Zool. Wood. Gen. Conch. f. 4.

Ch. cinereus, Zool. Jour. v. 5. p. 99. Wood Gen. Conch. f. 5. Chemn. Conch. 96. f. 818?

Red var. *C. ruber*? Chemn. Conch. t. 96. f. 813.

Black var. *C. Asellus*, Chemn. Conch. t. 96. f. 816.

Banded var. *C. minimus*, Chemn. Conch. t. 96. f. 814

C. marginatus and varieties, Sow. Jun. Conch. Illustr. f. 106 to 112.

Obs. This common species, including many varieties in colour, is more clearly referrible to *C. Marginatus*, of Pennant, than to *C. cinereus*, of authors; and without presuming to decide whether the description of Linnæus and the figure of Chemnitz were taken from ash coloured varieties of this or of the following species, I prefer applying the respective names as used by collectors at present. *C. marginatus* is more angular than the next species, and the granulations on the dorsal areas are not arranged in longitudinal lines.

C. minimus and *C. asellus*, of Chemnitz, may be black and banded varieties of this or of the next.

Sp. 4. *C. CINEREUS*.

Descr. Shell depressed, sub-carinated, minutely granulated; granulations on the central areas arranged in lines; margin minutely grained; colour various.

Ch. cinereus, Linn. Syst. Nat. p. 1106?

White var. *C. albus*, Auct?

Striped, *C. cimex*, Chemn. Conch. t. 96. 815.

Black, *C. Asellus*, Zool. Journ. vol. 5. pl. 5. f. 4.

Banded. *C. minimus*, Chemn. Conch? *Asellus*. (Young) Zool. Journ. v. 5. pl. 5. f. 3.

Ch. cinereus and varieties, Sow. Jun. Conch Illustr. Chiton, f. 94 to 98.

Obs. The comparative roundness and flatness of the shell, and the linear arrangement of the dorsal granules in this species distinguish it from the preceding. The white variety may have been the *C. albus*, of authors, but there is no proof of it, and having little else than colour to guide us, in reference to the ancient figures, it cannot be determined whether *C. Asellus*, of Chemn. is my black variety of *C. marginatus*, or Mr. Lowe's black var. of this species. The same difficulty occurs with respect to all the varieties.

Sp. 5. *C. CANCELLATUS*. Leach?

Descr. Shell elevated, narrow, with granulated striæ on the dorsal and lateral areas; margin minutely granular.

C. cancellatus, Sow. Jun. Conch. Illustr. f. 104. 104.
a. b. 105.

Obs. This minute species is narrower, more elevated, and more distinctly striated, than the preceding.

Sp. 6. *C. ALBUS*.

Descr. Shell oval, depressed, sub-carinated, lightly and minutely granulated; margin scaly.

C. albus, Linn. Syst. Nat. ? Mont. test. Brit.

Brown, var. *C. Aselloides*, Lowe, Zool. Journ. v. pl. v. f. 5.

C. albus and var. Sow. Jun. Conch. Illustr. Chiton 99. 99. a. 100.

Obs. Mr. Lowe, in a subsequent communication, states that his *C. Aselloides* is a variety of the *C. albus*, of Montagu. The latter may or may not be the *C. albus*, of Linnæus. It differs from the preceding in the arrangement of the granules, not being linear, and from *C. marginatus* in the large size, and regular arrangement of the granules, or rather scales of the marginal integument.

Sp. 7. *C. LÆVIS*.

Descr. Shell apparently smooth, but finely granulated, narrow, carinated: margin broad, velvety to the eye, but finely reticulated.

C. lævis, Pennant, Brit. Zool. 4. t. 36. f. 3? Lowe, Zool. Journ. vol. 5. p. 98. pl. 5. f. 1.

C. marginatus, Linn. Trans. viii. pl. 1. f. 2.

C. lævis and var. Sow. Jun. Conch. Illustr. Chiton
f. 101. 101. a. b. 102.

Sp. 8. *C. RUBER*.

Descr. Shell smooth, shining; valves beaked; margin minutely granular.

C. ruber, Linn. Syst. Nat.? Lowe, Zool. Journ. vol. 5. p. 101. pl. 5. f. 2. Sow. Jun. Conch. Illustr. Chiton, 103, 104.

Obs. Care must be taken to distinguish this from the red variety of *C. marginatus*, which it resembles in appearance, and to the unassisted eye seems almost equally smooth.

Sp. 9. *C. LATUS*.

Descr. Shell broad, smooth, sub-carinated, margin smooth; colour white and red, variegated.

C. latus, Lowe, Zool. Journ. vol. 5. pl. 5. f. 6, 7. Sow. Jun. Conch. Illustr. Chiton, f. 113, 113. a.



THE FOLLOWING BRITISH CHITONES ARE CONTAINED IN
THE
CONCHOLOGICAL ILLUSTRATIONS,

Parts 165, 166, 167, 168.

- Fig. 87. *Chiton fascicularis*, two views, enlarged one third. *a*, 5th. valve, magnified.
88 to 93. *C. crinitus*. 88. *a*, 5th valve, magnified.
92, Specimen with the hairs rubbed off, as frequently seen.
94 *C. cinereus*; *a*, fifth valve magnified.
95 to 98. *C. cinereus* and varieties. 98, *a*, 5th valve of white variety, magnified.
99. *C. albus*, enlarged. *a*, 5th valve magnified.
100. *C. albus*, var. *Aselloides*, *Lowe*.
101, 102. *C. lævis*. 102. *a*, enlarged. 101. *a*, 5th valve magnified. *b*, texture of margin.
103. *C. ruber*. *a*, inside; *b*, 5th valve, magnified.
104. *C. cancellatus*. *a*, 5th valve mag. *b*, figure enlarged.
105. *cancellatus* var.
106 to 112. *C. marginatus*, and varieties. 106, *a*, 5th valve, magnified; 110 *a*, 5th valve of red variety, magnified.
113. *C. latus*, *Lowe*, 2 views.

THE

THE

CONCHOLOGICAL ILLUSTRATIONS

PLATE 103, 104, 105

- Fig. 87. *Cyprina* *testudinaria*, two views, enlarged one
third, a. 5th valve, magnified.
88 to 93. *C. testudinaria*, 88, a. 5th valve, magnified.
89. Specimen with the 5th valve rubbed off, as
frequently happens.
90. *C. testudinaria*; a. 5th valve magnified.
91 to 93. *C. testudinaria* and varieties, 91, a. 5th
valve of white variety, magnified.
92. *C. testudinaria*, a. 5th valve magnified.
100. *C. testudinaria*, var. *Arctostoma*, Lamar.
101, 102. *C. testudinaria*, 101, a. 5th
valve magnified, 102, a. 5th valve of margin.
103. *C. testudinaria*, a. 5th valve; 104, 5th valve, magnified.
104. *C. testudinaria*, a. 5th valve, var. *Arctostoma*, Lamar.
105. *C. testudinaria*, var. *Arctostoma*, Lamar.
106 to 112. *C. testudinaria* and varieties, 106, a.
5th valve, magnified; 110 a. 5th valve of red
variety, magnified.
113. *C. testudinaria*, Lamar, 2 views.

Chiton.



1



2



3



4



5



6



7



8

Chiton.

10



9



11



12



13*



14



13

G.B.S. jun.



Chiton.

15



16



17



18*



19



18



20



21



Chiton.

22



23



24



25



26



27



28



29



30

Chiton.



32



31




33



34



35



36



37

Chiton.

38



39



40



41



42



43

Chiton.

44*



45



46



44



47



48



44**

Chiton.

49



50



51




52



53




54



55

Chelon



Chelon

Chiton



56



63



a



57



59



58




61



60



62



Chiton



64



67



67



67



65



67




67



68



a



Chiton



1841



Chiton



79



75



77



72



69



76





78



74



76



Chiton

83



83a



82



80



81



81



Chiton



86



84.d



84



80



85



85

Chiton



Chiton



87. a



90



89



88



93



88. a



92



91

Chiton



Chiton



87



99



87



98



99.a



97



100



98.a



96



95



94.a



94

Chiton



11



12




13



14



15



16




17




18



19



20




21



22

Chiton



Chiton



106



107



108



113



106.a



109



110.a



110



110



111



113.a



112

Chiton.



117a.



118



114



116



115



112



117



117

Carlin



Chiton



123



124



125



119



122



123



120



121



121



125



126



127




128



129



130



131



132

Chiton



126



128 n



127



126



130



128




129




128

Chiton



Chiton



Chiton

135



137



135



136




139



138




139



Chiton





Chiton

151



148



154




147



152

Chiton



Chiton



147



150



153



155



149



147



148

Chiron



14



15



16



17



18



19



20

A CATALOGUE
OF THE
RECENT SPECIES OF FISSURELLA,
BY G. B. SOWERBY.

- 1 *Fissurella picta*, *Lam.*, Conch. Illust. f. 4, *var.* f. 26. Terra del Fuego.
- 2 ——— *nimbosa*, *Lam.*, Conch. Illust. f. 2. West Indies.
- 3 ——— *crassa*, *Lam.* Conch. Illust. f. 9, 11, *F. Clypeiformis* mihi, Tank. Cat. app. p. vi. Lamarck's *F. depressa* is only a worn fragment of this species, as Mr. Gray informs me. Valparaiso.
- 4 ——— *græca*, *Lam.* Conch. Illust. f. 3. Nevis and the West Indian Islands generally.
- 5 ——— *nodosa*, *Lam.* Conch. Illust. f. 1. Trinidad, Antigua St. Vincents, &c.
- 6 ——— *Cayenensis*, *Lam.* Conch. Illust. f. 7. Monte Christi, West Columbia, Mr. Cuming. Obs. In G. Humphrey's collection were many specimens of this, but without locality.
- 7 ——— *lilacina*, *Lam.* Obs. This is entirely unknown to me.
- 8 ——— *rosea*, *Lam.* Conch. Illust. f. 8, West Indies.
- 9 *Barbadensis*, *Lam.* Conch. Illust. f. 5. *var.* f. 6, *var. alter.* f. 76, Barbadoes and other Islands of the West Indies.
Obs. The *var.* represented f. 76 has been considered as a distinct species, and named *virescens* by Mr. Guilding.

- 10 *Fissurella radiata*, *Lam. Conch. Illust.* f. 62, var. f. 64, West Indies

Obs. This is perhaps only a variety of *F. Barbadosensis*, it differs however from that species in the form of the dorsal aperture.

- 11 — *viridula*, *Lam. Conch. Illust.* f. 13, Trinidad, Antigua, &c.

- 12 — *hiantula*, *Lam. Conch. Illust.* f. 10, Southern Africa.

Obs. For a representation of this species *Lamarck* refers to *Born. Vign.* f. F. which is the same as I named *F. aperta* in the Tankerville Catalogue. The following are therefore synonyms of this species, viz: *Fissurella aperta*, *Tank. Cat. app. p. vi.* *Pupillæa aperta*, *Gray* in *Supp. to Beechey's Narrative*

- 13 — *Pustula*, *Lam. Conch. Illust.* f. 20, Sowerby Genera, No. xxi. f. 3, Trinidad, Antigua, &c.

- 14 — *fascicularis*, *Lam. Conch. Illust.* f. 14, Sowerby Genera, f. 6, West Indies.

- 15 — *Javanicensis*, *Lam. Conch. Illust.* f. 12, Cape of Good Hope.

- 16 — *Peruviana*, *Lam.* This is entirely unknown to me.

- 17 — *gibberula*, *Lam. Conch. Illust.* f. 17, Panama, Mr. Cuming.

- 18 — *minuta*, *Lam. Conch. Illust.* f. 16, Mediterranean.

- 19 — *crenulata, mihi*, *Tank. Cat. app. p. vi. Conch. Illust.* f. 31, 38.

Obs. I have only seen four specimens of this species, they are in the collections of Dr. Goodall, the Manchester Museum, Mr. Cuming, and Mr. Lincoln; it is from the specimen of the last named gentleman, as being the best

of the four, that our figures were drawn. I am informed by that gentleman that it was found at Monterey, in Upper California.

- 20 *Fissurella calyculata, mihi*, Genera of Shells, No. xxi. f. 4. Conch. Illust. f. 19, 21.
- 21 ——— *maxima, mihi*, Zool. Proc.* p. 123. Conch. Illust. f. 18, Valparaiso, Mr. Cuming.
- 22 ——— *grandis, mihi*, Zool. Proc. p. 123. Conch. Illust. f. 48, Valparaiso and Chiloe, Mr. Cuming.
- 23 ——— *limbata, mihi*, Zool. Proc. p. 123. Conch. Illust. f. 66, 74. var. f. 42, Syn. *F. picta, mihi*, Genera f. 1, Valparaiso, Mr. Cuming.
- 24 ——— *biradiata Frembl.* M. S. Zool. Proc. p. 124. Conch. Illust. f. 23. Valparaiso, Mr. Cuming. var. Conch. Illust. f. 52, Iquiqui, Mr. Cuming.
- 25 ——— *lata mihi*, Zool. Proc. p. 124. Conch. Illust. f. 63, † St. Mary's Island, Chili, Mr. Cuming.
- 26 ——— *pulchra, mihi*, Zool. Proc. p. 124. Conch. Illust. f. 24, Valparaiso, Mr. Cuming.
- 27 ——— *oriens, mihi*, Zool. Proc. p. 124. Conch. Illust. f. 25, var. f. 60, Island of Chiloe, Mr. Cuming.
- 28 ——— *Chilensis, mihi*, Zool. Proc. p. 124. Conch. Illust. f. 36, Valparaiso, Mr. Cuming.
- 29 ——— *obscura, mihi*, Zool. Proc. p. 125. Conch. Illust. f. 27, Gallapagos Islands, Mr. Cuming.
- 30 ——— *coarctata, King*, Zoological Journal, vol, 5. p. 339. Conch. Illust. f. 22, Sowerby Genera of Shells, No. xxi. f. 2 (*Fissurella crassa*). Port Praya.

* Proceedings of the Zoological Society of London, 1834.

† Miscalled *F. Chilensis* in the list published with the 76th and 77th parts. It is perhaps as well here to observe that these said lists are all to be cancelled eventually.

- 31 *Fissurella virescens, mihi*, Zool. Proc. p. 125. Conch. Illust. f. 37, * Panama, Mr. Cuming.
- 32 — *nigro-punctata, mihi*, Zool. Proc. p. 125. Conch. Illust. f. 51, Gallapagos and Lobos Island, Mr. Cuming.
- 33 — *macrotrema, mihi*, Zool. Proc. p. 125. Conch. Illust. f. 41, Gallapagos and Real Llejos, Mr. Cuming. var. Conch. Illust. f. 41, * Lobos Island, Mr. Cuming. var. Conch. Illust. f. 41, ** Lambeyeque, Peru, Mr. Cuming.
- 34 — *hians, Broderip and Sowerby*, Zool. Journ. iv. p. 370, †
- 35 — *affinis, Gray, Supp. to Beechey's Narrative*, Zool. Proc. p. 125. Conch. Illust. f. 44, Coast of Peru, Mr. Cuming. var. Valparaiso, Mr. Cuming.
- 36 — *inaequalis, mihi*, Zool. Proc. p. 126. Conch. Illust. f. 45, Guacomayo and the Gallapagos Islands, Mr. Cuming.
- 37 — *Pica, mihi*, Zool. Proc. p. 126. Conch. Illust. f. 32, St. Elena, Mr. Cuming. Variety, Conch. Illust. f. 33, Monte Christi, Mr. Cuming.
- 38 — *cancellata, mihi*, Tank. Cat. No. 802. Conch. Illust. f. 29. *Patella cancellata*, Sol. M. S.; *Patella ventricosa* Gmel. according to Mr. Gray; *Feraminella Sowerbii*, Guilding M. S.; *Lucapina elegans* Gray, St. Vincents.
- 39 — *microtrema, mihi*, Zool. Proc. p. 125. Conch. Illust. f. 50, Real Llejos, Mr. Cuming.
- 40 — *costata, Gray*. Shell elliptical, rather depressed, with a few radiating obtuse ribs, externally greenish,

* Miscalled *F. coarctata* King, in the published list.

† The specimen being mislaid, this species cannot be figured.

smooth; within white with an undulated margin: dorsal aperture very small, elliptical; internal margin broad: length 2.0, breadth 1.5 inches. Conch. Illust. f. 28, St. Jago.

- 41 *Fissurella olivacea*, Gray in Brit. Mus. Conch. Illust. f. 57.

- 42 ——— *Scutellum*, Gray in Brit. Mus. Conch. Illust. f. 34, Cape of Good Hope.

Obs. This is *F. Trapezina mihi*, in Zool. Proc. p. 126.

- 43 ——— *Europæa, mihi*, Conch. Illust. f. 43. Shores of the British Channel, &c.

Obs. This is the *Patella græca* of British Conchological writers, and the *F. cancellata* of Gray in Annals of Philos. new series vol. 1. The *Patella apertura* of Montagu is the young of it.

- 44 ——— *Mediterranea*, Gray in Brit. Mus. Conch. Illust. f. 30, Mediterranean.

- 45 ——— *macroschisma* Humphrey, Conchology. Conch. Illust. f. 39, New South Wales. var. f. 39,* Swan River.

Obs. This forms the Genus *Macroschisma* of Gray.

- 46 ——— *Noachina mihi*, Conch. Illust. f. 15, Leach's British Shells, Tab. x, f. 4, 5 (*Cemoria Flemingii*).

Obs. This is *Patella Fissurella*, Müll. Zool. Dan. t. 24, f. 4—6. *Patella Noachina*, Chemn, t. 197, f. 1927, 1928, who says it is British; and the Genus *Puncturella* of Lowe, Zool. Journ. iii. p. 77, who found it at Oban in Argyleshire.

- 47 ——— *Chemnitzii mihi*, Conch. Illust. f. 55. Zool. Proc. p. 126.

- 48 *Fissurella latimarginata, mihi*, Zool. Proc. p. 126. Conch. Illust. f. 69, Valparaiso and Iquique, Mr. Cuming.
- 49 — *æqualis mihi*, Zool. Proc. p. 127. Conch. Illust. f. 56, St. Elena, Mr. Cuming.
- 50 — *fulvescens, mihi*, Zool. Proc. p. 127. Conch. Illust. f. 49, Valparaiso, Mr. Cuming.
- 51 — *nigrita, mihi*, Zool. Proc. p. 127. Conch. Illust. f. 47, Cape of Good Hope.
- 52 — *aspera, mihi*, Zool. Proc. p. 127. Conch. Illust. f. 46, Pacosmayo, Peru, Mr. Cuming.
- 53 — *asperella, mihi*, Zool. Proc. p. 127. Conch. Illust. f. 71, Lobos Island, Peru, Mr. Cuming.
- 54 — *mutabilis, mihi*, Zool. Proc. p. 127. Conch. Illust. f. 67, 70, Cape of Good Hope, Mr. Cuming. Brazil? Humphrey
- 55 — *Panamensis, mihi*, Zool. Proc. p. 127. Conch. Illust. f. 72, var. f. 72,* Panama, Mr. Cuming.
- 56 — *Ruppellii, mihi*, Zool. Proc. p. 128. Conch. Illust. f. 65, 75, Nevis, Mr. Powers; Red Sea, Rüppell.
- 57 — *Clypeus, mihi*, Zool. Proc. p. 128. Conch. Illust. f. 77, St. Elena, Mr. Cuming.
- 58 — *subrostrata, Gray,** Conch. Illust. f. 35. Shell oval, somewhat depressed, anteriorly subrostrated, extremities raised; white within, with a crenated margin; externally radiately ribbed, ribs rather distant, muricated; dorsal aperture small, elliptical, its internal edge broad: length 1·1, breadth 0·8 inches. Mr. Gray has a single specimen in his own collection, from St. Vincents.
- 59 — *sanguinea, Gray*, Conch. Illust. f. 53. Shell ovate-oblong, depressed, externally white, radiately ribbed

* Not Guilding, as printed incorrectly in the list.

and striated; ribs obtuse, their interstices spotted with bright red; inside white, somewhat transparent, edge crenated; dorsal aperture oblong.

Obs. I have only seen one specimen which is in Mr. Gray's Collection.

- 60 *Fissurella Tongana*, *Quoy and Gaimard*, Zool. de l'Astrolabe III. p. 335. t. 68. f. 3. 4. Tonga-tabou.

Obs. I have never seen this species.

- 61 — *Afra*, *Quoy and Gaimard*, Zool. de l'Astrolabe III. p. 336. Tab. 68. f. 5. 6. Cape de Verd Islands.

Obs. I have never seen this species.

- 62 — *Lincolni*, *Gray*. Conch. Illust. f. 40. Monterey, Upper California.

Obs. A splendid species obligingly lent by Ab. Lincoln, Esq. to be drawn for this work. The only specimen hitherto known is in his collection.

- 63 — *candida, mihi*, Conch. Illust. f. 58. Shell oblong, white, subhyaline, externally marked with distinct decussating striæ, dorsal aperture anterior, oblong, wider in the middle, contracted at each end, its internal margin broad, truncated posteriorly. Locality unknown. In my own and Mr. Cuming's collection.

- 64 — *obtusa mihi*, Conch. Illust. f. 59. Cape of Good Hope, Humphrey. Shell elliptical, rather elevated, pale coloured, with radiating striæ and dark purplish bands; inside green; dorsal aperture elliptical; its internal margin broad, equal, sometimes with a dark purple edge.

Obs. All the specimens I have seen were in G. Humphrey's collection.

- 65 — *lineata mihi*, Conch. Illust. f. 68. Shell oblong,

rather elevated and obtuse at both extremities; cancellately ribbed; a few of the ribs pale purple; dorsal aperture small, much nearer to the anterior than to the posterior end, its margin internally truncated posteriorly: margin crenated.

Obs. The only two specimens I have seen were in G. Humphrey's collection, they are now in Mr. Cuming's.

66 *Fissurella rugosa, mihi*. Conch. Illust. f. 54. Shell of a rather lengthened oval shape, depressed, whitish, varied with dark red brown, and covered with a few large rugose radiating ribs and some smaller interstitial ones; dorsal aperture nearer to the anterior extremity, oblong, its internal margin rather wide, with a dark border; inside green; edge of the whole shell irregularly sinuous: length 1.0, breadth 0.6 inches, Gallapagos Islands, Mr. Cuming.

67 — *Mexicana mihi*, Conch. Illust. f. 61. Shell oblong, reddish brown, with darker rays, and numerous close-set radiating grooves, which are decussated by very fine concentric striæ; dorsal aperture oblong, rather contracted in the middle, and with two obsolete teeth on each side; inside white, with a purplish brown crenulated margin: length 1.25, breadth 0.56. Real Llejos, Mexico; Mr. Cuming.

68 — *crenifera, mihi*, Zool. Proc. p. 128. Conch. Illust. f. 73, Real Llejos, Mr. Cuming.

LIST OF FIGURES OF FISSURELLÆ

In Parts 68 to 78, and 80 of the


Conchological Illustrations.

- Fig.
1 *F. nodosa* Lam.
2 *F. nimbosa* Lam.
3 *F. Græca* Lam.
4 *F. picta* Lam.
5 *F. Barbadenis* Lam.
6 *F. ———* var.
7 *F. Cayenensis* Lam.
8 *F. rosea* Lam.
9 *F. crassa* Lam. young.
10 *F. hiantula* Lam.
11 *F. crassa* Lam.
12 *F. Javanicensis* Lam.
13 *F. viridula* Lam.
14 *F. fascicularis* Lam.
15 *F. Noachina* nobis.
16 *F. minuta* Lam.
17 *F. gibberula* Lam.
18 *F. maxima* nobis.
19 *F. calyculata* nobis.
20 *F. Pustula* Lam.
21 *F. calyculata* nobis.
22 *F. coarctata* King.
23 *F. biradiata* Frembly.
24 *F. pulchra* nobis.
25 *F. Oriens* nobis.
26 *F. picta* Lam. var.
27 *F. obscura* nobis.
28 *F. costata* Gray.
29 *F. cancellata* nobis.
30 *F. mediterranea* Gray.
31 *F. crenulata* nobis.
32 *F. Pica* nobis.
33 *F. Pica* var. nobis.
34 *F. Scutellum* Gray.
35 *F. subrostrata* Guilding.

- 36 *F. Chilensis nobis.*
- 37 *F. coarctata King. virescens Saw.*
- 38 *F. crenulata nobis.*
- 39 *F. macroschisma.*
- 39* *F. macroschisma var.*
- 40 *F. Lincolni Gray.*
- 41 *F. macrotrema nobis. 41* F.*
- 41** *F. macrotrema var.*
- 42 *F. limbata var. nobis.*
- 43 *F. europæa nobis.*
- 44 *F. affinis Gray.*
- 45 *F. inæqualis nobis.*
- 46 *F. asper^a nobis.*
- 47 *F. nigrita nobis.*
- 48 *F. grandis nobis.*
- 49 *F. fulvescens nobis.*
- 50 *F. microtrema nobis.*
- 51 *F. nigropunctata nobis.*
- 52 *F. biradiata var.*
- 53 *F. sanguinea nobis.*
- 54 *F. rugosa nobis.*
- 55 *F. Chemnitzii nobis.*
- 56 *F. æqualis nobis.*
- 57 *F. olivacea Gray.*
- 58 *F. candida nobis.*
- 59 *F. obtusa nobis.*
- 60 *F. oriens var. nobis.*
- 61 *F. Mexicana nobis.*
- 62 *F. radiata Lam.*
- 63 *F. Chilensis nobis.*
- 64 *F. radiata Lam. var.*
- 65 *F. Ruppellii nobis.*
- 66 *F. limbata nobis.*
- 67 *F. mutabilis nobis.*
- 68 *F. lineata nobis.*
- 69 *F. latimarginata nobis.*
- 70 *F. mutabilis var.*
- 71 *F. asperella nobis.*
- 72 *F. Panamensis nobis.*
- 72* *F. Panamensis var.*
- 73 *F. crenifera nobis.*
- 74 *F. limbata nobis.*
- 75 *F. Ruppellii nobis. Red Sea.*
- 76 *F. Barbadosensis var.*
- 77 *F. Clypeus nobis.*

Fissurella.






Fissurella.

9



8



10



11



12



13



14

Fissurella.




16



15



17



18



19



20



21

G.B.S. jun. etc.

Fissurella.

23



22




23



24



25



26



27


G.B.S. jun. exc.

Parurella




Fissurella.


29




29




28



30



30



31



32



33

G.B.S.

Fissurella.



34



35




36



37



38



39




40



41

G.B.S. jun. sc.



Fissurella.




41*



41



41**



40



42



43



44

Fissurella.




45



46




47




48



49



50



51

GBS. jun. 1800.




100



101



102



103



104



105



106

Fissurella.


54



53



54




55



56




57



52



Fissurella.



G.B. Sjun. exc.

Fig. 1




Fissurella.

64



65




66




67




68




69






Monoceros.



G.B.S. jun. exc.



Fissurella.

70



71



72



73



74



72*



75



776



77

G.B.S. for exc

Sporella



BULINUS.

LIST OF FIGURES,

In Parts, 21, 22, 23, 26, 27, 30, 31, 34, 35, 137 to 146, 185, 186, of the CONCHOLOGICAL ILLUSTRATIONS,

BY

G. B. SOWERBY, Jun.

- Fig. 1*B. Broderipii, *Sow. Z. P.* 1832. Copiapo in Chili.
1**B. Broderipii var. *Sow. Z. P.* 1832. Iniqui in Peru.
2 B. nitidus *Brod. Z. B.* 1832. Tumbez in Peru.
2* B. Cattivorus *Brod. Z. P.* 1832. Monte Christi in West Columbia.
3 B. Coturnix *Sowerby, Z. P.* 1832. Huasco in Chili.
4 B. Chilensis *Sowerby, Coquimbo.*
5 B. rosaceus *King, Zool. Journ. vol. v. p. 341, Valparaiso.*
6 & 6*B. vittatus *Brod. Z. P.* 1832, Ilo, Peru.
7 B. granulatus *Brod. Z. P.* 1832, Valparaiso.
7* B. granulatus var. *Brod. Z. P.* 1832. Conception.
8 B. Coquimbensis *Brod. Z. P.* 1832, Coquimbo.
9 *Partula hyalina, Brod. Z. P.* 1832, Oheataroa or Rurutu
10 B. guttatus *Brod. Z. P.* 1832, Cobija, Peru.
11 B. translucens *Brod. Z. P.* 1832, Panama
12 B. Gravesii *King, Zool. Journ. v. p. 340, Maule, Chili.*
12* B. Gravesii var. Conception.
12**B. Gravesii, var. Quilliotta.
13 B. Scalariformis *Brod. Z. P.* 1832, Ancon, Peru.

- 14 B. *Proteus* *Brod.* Z. P. 1832. Truxillo.
- 14a B. *Proteus* var. Z. P. 1832. St. Jacinta.
- 14b B. *Proteus* var. *Brod.* Z. P. 1832. Pacosmayo.
- 14c B. *Proteus* var. *Brod.* Z. P. 1832. Lima.
- 15 B. *mutabilis* *Brod.* Z. P. 1832. Santos, Peru.
- 15* B. *mutabilis* var. *Brod.* Z. P. 1832. Campania
of Truxillo.
- 16 B. *versicolor* *Brod.* Z. P. 1832. Mongon, near
Casma.
- 16* B. *versicolor* var. *Brod.* Z. P. 1832. Mongon,
near Casma.
- 17 B. *pulchellus* *Brod.* Z. P. 1832. Truxillo.
- 18 B. *Princeps* *Brod.* Z. P. 1832. Conchagua, Cent-
ral America.
- 19 B. *modestus* *Brod.* Z. P. 1832. Huacho, Peru.
- 20 B. *varians* *Brod.* Z. P. 1832. Truxillo, Peru.
- 20* B. *varians* var. *Brod.* Z. P. 1832.
- 21 B. *Tigris* *Brod.* Z. P. 1832. Truxillo, Peru.
- 22 B. *albicans* *Brod.* Z. P. 1832. Huasco, Chili.
- 22* B. *albicans* var. *Brod.* Z. P. 1832. Copiapo, Chili.
- 23 B. *pustulosus* *Brod.* Z. P. 1832. Huasco.
- 24 B. *scabiosus* *Sow.*
- 25 B. *Panamensis* *Brod.* Z. P. 1833. I. of Saboga,
Panama.
- 26 B. *vexillum* *Brod.* Z. P. 1832. Isle of Saboga.
- 27 B. *pupifornis* *Brod.* Z. P. 1832. Huasco.
- 28 B. *chrysalidiformis* *Sow.* Z. P. 1833.
- 29 B. *bilineatus* *Sow.* Z. P. 1833. St. Elena.
- 30 B. *affinis* *Brod.* Z. P. 1832. Hill of Mexillones.
- 31 B. *turritus* *Brod.* Z. P. 1832. Truxillo.
- 32 B. *rubellus* *Brod.* Z. P. 1832. Truxillo.
- 33 B. *Nucleus* *Sow.* Straits of Magalhæns.
- 34 B. *arrosus* *Brod.* Z. P. 1832.
- 35 B. *Javanicus* *Sow.* }
- 35* B. do. young } Java.
- 36 B. *punctulifer* *Sow.* Z. P. 1833. Questo Prado,
Chili.
- 37 B. *Nux.* *Brod.* Z. P. 1832. Galapagos Islands.
- 37* B. *Nux* *Brod.* Z. P. 1832. Galapagos Islands.
- 38 B. *derelictus* *Brod.* Z. P. 1832. Cobija.
- 39 B. *scutulatus* *Brod.* Z. P. 1832. Islay, Peru.
- 39* B. *pallidior, nobis.* South Sea Islands.
- 40 B. *rugiferus nobis.* Gallapagos Islands.
- 41 B. *calvus nobis.* Gallapagos Islands.
- 42 B. *ustulatus nobis.* Gallapagos Islands.
- 43 B. *unicolor, nobis.* Panama.
- 44 B. *pallidior, nobis.* var.

- 45 B. Jacobi, *nobis*, two varieties.
James's Island, one of the Gallapagos.
- 46 B. bivittatus, *nobis*. Brazil.
- 47 B. opalinus, *nobis*. Brazil.
- 48 B. laticolor, *nobis*.
- 49 B. conspersus *nobis*. Lima.
- 50 B. erythrostoma *nob.* Copiapo.
- 51 B. albus *Brod.* Copiapo.
- 52 B. discrepans *nobis*. San Salvador.
- 53 B. Luzonicus *nob.* ovoideus Brug. Manilla.
- 54 B. undulatus. *Bulinulus undulatus*. Guilding, St.
Vincent.
- 55 B. unifasciatus *nob.* Galapagos Islands.
- 56 B. striatus *King*. Santos, Peru.
- 57 B. signatus. *Macroceramus signatus* Guilding, St.
Vincent.
- 58 B. striatulus *nobis*. Hills of Lima.
- 59 B. Mahogani *nobis*.
- 60 B. Crichtoni *Brod.* two views.
- 61 B. inflatus *Brod.* three views.
- 62 B. hygrophylæus *D'Orb.* two views.
- 63 B. maximus *Sowerby*, in *Tank. Cat.*
Helix kremnoica D'Orb. one view.
- 64 B. apodemetes *D'Orb.* two views.
- 64* B. apodemetes *var.* one view.
- 65 B. lophoicus *D'Orb.* Bolivia.
- 66 B. Pusio *Brod.*
- 67 B. Inca *D'Orb.*
- 68 B. pæcilus *D'Orb.*
- 69 B. pæcilus *var.* *D'Orb.*
- 70 B. thamnoicus *var.* *D'Orb.*
- 71 B. thamnoicus *var.* *D'Orb.* two views.
- 72 B. thamnoicus *var.* *D'Orb.*
- 73 B. Sporadicus *D'Orb.* Corrientes.
- 74 B. Sporadicus *D'Orb.* Corrientes.
- 75 B. Sporadicus *D'Orb.* Corrientes.
- 76 B. auritus. *nob.* Bahia.
- 77 B. Tupacii *D'Orb.*
- 78 B. Sporadicus *D'Orb.* Bolivia.
- 79 B. melanostoma *var?*
- 80 B. Onca *D'Orb.*
- 81 B. Lychnorum *D'Orb.*
- 82 B. Lychnorum *var.* alba. *D'Orb.*
- 83 B. Bacterionides *D'Orb.*
- 84 B. montivagus *D'Orb.*
- 85 B. eschariferus *nob.* Galapagos.
- 86 B. culmineus *D'Orb.*

- 87 *B. rugulosus* *nob.* Galapagos.
 88 *B. melanostoma* var.
 89 *B. multicolor*.
 90 *B. Myersii*, *Miller*.
 91 *B. pulchellus*, *nob.* Huallaga.
 92 *B. pulchellus* var. Huallaga.
 93 *B. piperitus*, *nob.* Huallaga.
 94 *B. Milleri*, *nob.*
 95 *B. strigatus*, *nob.* Huallaga.
 96 *B. strigatus* var. Huallaga.
 97 *B. Janeirensis*, *nob.* Rio de Janeiro.
 98 *B. perplexus* *nob.*
 99 *B. Downesii* *Gray*.
 100 *B. maculiferus* *nob.*
 From Lady Harvey's Collection.
 101 *B. hæmastoma* var.
 102 *B. proximus* *nob.*
 103 *B. hæmastoma*, var.
 104 *B. ovoideus*, *Brug.*
 105, 106, & 107 *B. ovoideus*, varieties.
 108 *B. porraceus*, *Jay*.
 109, 110, & 111 *B. porraceus*, varieties.
 112 *B. virgatus*, *Jay*.
 113 *B. virgatus*, var.
 114 *B. virgatus*, var.
 115 *B. fictilis*, *Brod.*
 116 *E. fictilis*, *Brod.* var.
 117 *B. larvatus*, *Brod.*

Bulinus.

- 107 *H. ...*
- 108 *H. ...*
- 109 *H. ...*
- 110 *H. ...*
- 111 *H. ...*
- 112 *H. ...*
- 113 *H. ...*
- 114 *H. ...*
- 115 *H. ...*
- 116 *H. ...*
- 117 *H. ...*
- 118 *H. ...*
- 119 *H. ...*
- 120 *H. ...*
- 121 *H. ...*
- 122 *H. ...*
- 123 *H. ...*
- 124 *H. ...*
- 125 *H. ...*
- 126 *H. ...*
- 127 *H. ...*
- 128 *H. ...*
- 129 *H. ...*
- 130 *H. ...*
- 131 *H. ...*
- 132 *H. ...*
- 133 *H. ...*
- 134 *H. ...*
- 135 *H. ...*
- 136 *H. ...*
- 137 *H. ...*
- 138 *H. ...*
- 139 *H. ...*
- 140 *H. ...*
- 141 *H. ...*
- 142 *H. ...*
- 143 *H. ...*
- 144 *H. ...*
- 145 *H. ...*
- 146 *H. ...*
- 147 *H. ...*
- 148 *H. ...*
- 149 *H. ...*
- 150 *H. ...*
- 151 *H. ...*
- 152 *H. ...*
- 153 *H. ...*
- 154 *H. ...*
- 155 *H. ...*
- 156 *H. ...*
- 157 *H. ...*
- 158 *H. ...*
- 159 *H. ...*
- 160 *H. ...*
- 161 *H. ...*
- 162 *H. ...*
- 163 *H. ...*
- 164 *H. ...*
- 165 *H. ...*
- 166 *H. ...*
- 167 *H. ...*
- 168 *H. ...*
- 169 *H. ...*
- 170 *H. ...*
- 171 *H. ...*
- 172 *H. ...*
- 173 *H. ...*
- 174 *H. ...*
- 175 *H. ...*
- 176 *H. ...*
- 177 *H. ...*
- 178 *H. ...*
- 179 *H. ...*
- 180 *H. ...*
- 181 *H. ...*
- 182 *H. ...*
- 183 *H. ...*
- 184 *H. ...*
- 185 *H. ...*
- 186 *H. ...*
- 187 *H. ...*
- 188 *H. ...*
- 189 *H. ...*
- 190 *H. ...*
- 191 *H. ...*
- 192 *H. ...*
- 193 *H. ...*
- 194 *H. ...*
- 195 *H. ...*
- 196 *H. ...*
- 197 *H. ...*
- 198 *H. ...*
- 199 *H. ...*
- 200 *H. ...*



Bulinus





Bulinus





Bulinus.

15



14



15*



14a.



14b.



14c.



16



16*



17

Bulinus.



19



18



20



21.



20*



18



22



22*

G.B.S. jun. ex.

Bulinus.

23



24



25



26



27



28



29



30



32



31

Bulinus.

33



34



35



36



35*



36



37



37*



38

1841



1842

Bulinus.

40



39



41



42



44



43



45



45




46




48



47

G. B. S. jun. ex.



Bulinus.



49



50



51



52



53



54



55



56



58



57





Bulinus

Pl. 138.



64.



64.



64.



66.



66.



63.




67.



67.

G. B. Smith del.





68



72



69



67



73



71



70



75



73



74



76



77



77



78

Bulinus



83.



83



80



81



82



79





84



84

G. B. S. Jan. exc.

Butyrus



Butyrus

Bulinus



85.



85



80



86



86



79.



87



87

G. B. L. Junr. exc.

Bulimus



Bulinus.



95



95



92



91



91



88



96



96



Bulinus.



97



97



93



90



90



94



89



89



Bulinus

98



98



99




100



100





101




103



102

103



Bulinus



109



104



108



105



111



106



110



Bulinus

116



117



115



112



114



113



107



A CATALOGUE

Of all the Species hitherto known of the Genus Chilina, of Gray,
represented in the 135th and 136th parts of

Conchological Illustrations,

By G. B. SOWERBY, JUN.

- Spec. 1 *Chilina Dombeyana*. Conch. Illustr. f. 11.
2. ——— *ampullacea*. *Sowerby*. Malacological and Con-
chological Mag. part 2. Conch. Illustr. f. 3.
3. ——— *tenuis*. *Gray*. Conch. Illustr. f. 12.
4. ——— *fluctuosa*. *Gray*. Spicilegia Zoologica, t. 6.
f. 19. Conch. Illustr. f. 2.
5. ——— *fluviatilis*. *Gray*. In Mus. Brit. Conch.
Illustr. f. 5.
6. ——— *fluminea*. *Gray*. Spicil. Zool. p. 5. Conch.
Illustr. f. 7.
7. ——— *major*. *Sowerby*. Malacol. and Conchol. Mag.
part 2, Conch. Illustr. f. 10.
8. ——— *robustior*. *Sowerby*. Malacol. and Conchol.
Mag. part 2. Conch. Illustr. f. 1.
9. ——— *Parchappii* *D'Orb.* Synops. Conch. Illustr.
f. 8.
10. ——— *Tehuelcha*. *D'Orb.* M. S. Conch. Illustr. f. 9.
11. ——— *Puelcha*. *D'Orb.* M. S. Conch. Illustr. f. 13.
12. ——— *gibbosa*. *Sowerby*. Malacol. and Conchol. Mag.
part 2. Conch. Illustr. f. 4.
13. ——— *ovalis*. *Sowerby*. Malacol. and Conchol. Mag.
part 2. Conch. Illustr. f. 6.

A CATALOGUE
LIST OF FIGURES.

Fig. 1. *Chilina robustior*

2. ——— *fluctuosa*.

3. ——— *ampullacea*.

4. ——— *gibbosa*.

5. ——— *fluviatilis*.

6. ——— *ovalis*.

7. ——— *fluminea*.

8. ——— *Parchappii*.

9. ——— *Tehuelcha*.

10. ——— *major*.

11. ——— *Dombeyana*.

12. ——— *tenuis* and var.

13. ——— *Puelcha*.

Chilina.



6



1



7



1




2



2



3



4



4



5

G. B. Smith: exc.

Plum



1. Plum



2. Plum

3. Plum

4. Plum

5. Plum



6. Plum

7. Plum

8. Plum



9. Plum

Chilina



8



9



9



8



10



11



10



12



13



12

G.B.S. f. v. 1855

Chama



A CATALOGUE

OF ALL THE RECENT SPECIES HITHERTO KNOWN
OF THE GENUS

Neritina,

Figured in Parts 86, 87, 90, 91, 94 to 100, of the
CONCHOLOGICAL ILLUSTRATIONS,

BY

G. B. SOWERBY, *Junr.*

- Spec. 1. *N. pulligera* Lam. Hist. nat. des. Anim. sans vert.
Encycl. method t. 455. f. 1. Conch. Illustr. f.
26. Genera of Recent and Fossil Shells, *Neritina*,
f. 6. India.
2. *N. canalis* Sowerby Tank. Cat. No. 1094, app. p.
xi. Wood Ind. Test. Sup. *Nerita*, f. 17. Conch.
Illustr. f. 22. Tahiti.
3. *N. dubia* Lam. Chemn. V. f. 2019. 2020. Wood
Ind. test. t. 35. *Nerita*. f. 29. Conch. Illustr.
f. 28.
4. *N. Zebra* Lam. Encycl. method. t. 455. f. 3. Wood
Ind. test. t. 35, *Nerita* f. 18. Tahiti.
5. *N. Ziczac* Lam. Conch. Illustr. f. 41.
6. *N. punctulata* Encycl. method t. 455. f. 2. Conch.
Illustr. f. 21 Isle St. Vincents.
7. *N. gagates* Lam. Conch. Illustr. f. 29.
8. *N. lugubris* Lam. Conch. Illustr. f. 38.
9. *N. Corona*. Lam. 1. c. Sowerby Genera of Recent
and Fossil Shells, *Neritina* f. 3. Conch. Illustr.
f. 20. Mauritius.

- Sp .10. *N. spinosa* *Budgin* M. S. *Sowerby*, Tank. Cat. No. 1102 Conch. Illustr. f. 9. Tahiti.
11. *N. brevispinosa* *Lam.* Conch. Illustr. f. 8. 8. *a. b. c. d. e.* (*Spinosa* *Wood* Sup. t. 8. *Nerita* f. 12.) Timor and Haynam. *Humphrey*.
12. *N. crepidularia* *Lam.* Wood Sup. t. 8. *Nerita* f. 6. Conch. Illustr. f. 25. River Hoogly, Calcutta.
13. *N. auriculata* *Lam.* Encycl. method. t. 455. f. 6. Conch. Illustr. f. 17. Sumatra.
14. *N. Domingensis* *Lam.* Conch. Illustr. f. 42. St. Domingo (*Lamarck*.)
15. *N. fasciata* *Lam.* Encycl. method. t. 455. f. 5. Conch. Illustr. f. 35.
16. *N. lineolata* *Lam.* Chemn. ix. f. 1081. Encycl. method. t. 455. f. 4. Conch. Illustr. f. 37.
17. *N. semiconica* *Lam.* Chemn. ix. f. 1087. Wood Sup. t. 8. *Nerita* f. 15 India. Conch. Illus. f. 23.
18. *N. strigilata* *Lam.* Conch. Illustr. f. 4. var. f. 32. (*turrita* Chemn. ix. f. 1085.) Sumatra.
19. *N. Meleagris* *Lam.* Chemn. ix. t. 124. f. D. L. Conch. Illustr. f. 19. (3 var.) (*Nerita pulchella* *Wood*. Sup. t. 8. f. 18.) Brazil.
20. *N. virginea* *Lam.* Chemn. ix. t. 124. f. H. I. Conch. Illustr. f. 27. (six var.) W. Indies.
21. *N. fluviatilis* *Lam.* *Drap.* t. 1. f. 3. 4. Conch. Illustr. f. 33. Europe.
22. *N. viridis* *Lam.* Chemn. ix. f. 1081. 1082. Conch. Illustr. f. 24. (2 var.) West Indies. Antigua.
23. *N. Bætica* *Lam.* Conch. Illustr. f. 45. Spain.
24. *N. granosa* *Sowerby* Tank. Cat. No. 1115. Conch. Illustr. f. 6. (*papillosa* *Jay* Cat. of Recent Shells, pl. 4. f. 11. 11. 11.
25. *N. Oweniana* *Gray.* Wood Sup. t. 8. *Nerita* f. 8. Conch. Illustr. f. 15. Africa.

- Sp. 26. *N. cariosa* *Gray*. Wood Sup. t. 8. Nerita f. 9.
Conch. Illustr. f. 5. (In the list published
with the 91st. part of Conch. Illustr. this is
incorrectly named *N. Caffra* *Gray*.)
27. *N. Caffra* *Gray*. Woods Sup. t. 8. Nerita. f. 10.
Conch. Illustr. f. 51. Fernando Po.
28. *N. Smithii* *Gray*. Wood Sup. t. 8. Nerita f. 11.
Conch. Illustr. f. 36. River Hoogly, Calcutta.
29. *N. latissima* *Brod.* Zool. Proc. Part II. 1832 p. 201.
Conch. Illustr. f. 3. var. Conch. Illustr. f. 16.
Real Llejos.
30. *N. dilatata*. *Brod.* Zool. Proc. 1. c. Conch. Illustr.
f. 11. Tahiti.
31. *N. globosa* *Brod.* Zool. Proc. 1. c. Conch. Illustr. f.
12. (*Owenii* Wood Sup. t. 8. Nerita f. 16.)
Chiriqui.
32. *N. Nux.* *Brod.* Zool. Proc. 1. c. Conch. Illustr. f.
39. Tahiti.
33. *N. intermedia* *Sowerby*, Zool. Proc. loc. cit. Conch.
Illustr. f. 7. Bay of Montejo and Gulf of
Nocoiya.
34. *N. chlorostoma*, *Sowerby*, Zool. Proc. 1. c. Conch.
Illustr. f. 34. Tahiti.
35. *N. picta*, *Sowerby*, Zool. Proc. 1. c. Conch. Illustr
f. 1. Panama.
36. *N. reticulata* *Sowerby*, Zool. Proc. 1. c. Conch.
Illustr. f. 2. Lord Hood's Island.
37. *N. Morio*, *Sowerby*, Zool. Proc. 1. c. Conch. Illustr.
f. 40. Ducie's and Easter Islands.
38. *N. Faba*, *Sowerby*, Conch. Illustr. f. 10. Singapore.
39. *N. Afra*, *Sowerby*, Conch. Illustr. f. 13. Fernando
Po.
40. *N. piperina* *Chemn.* Conch. Illustr. f. 18. Malabar,
Chemn. Found among Pepper.
41. *N. subgranosa*, *Sowerby*, Conch. Illustr. f. 14.
Timor.

- Sp. 42. *N. Prevostiana* Conch. Illustr. f. 46. Europe.
 43. *N. Danubialis*, *Sadler*, Conch. Illustr. f. 47. Danube.
 44. *N. Dalmatica*, *Partsch*, Conch. Illustr. f. 57.
 Dalmatia.
 45. *N. Pupa*, *Dillw*, Conch. Illustr. f. 30. Providence
 Island, and Haynam.
 46. *N. lutescens*, *Muhlf.* Conch. Illustr. f. 56. (varia.
 Parr.) Cephalonia.
 47. *N. rivalis*, *Parreyss*, Conch. Illustr. f. 58. Europe.
 48. *N. Jordani*, *Sowerby*, Conch. Illustr. f. 49. (3 var)
 River Jordan, Dr. Butler.
 49. *N. labiosa*, *Sowerby*, Conch. Illustr. f. 48.
 50. *N. subsulcata*, *Sowerby*, Conch. Illustr. f. 50.
 51. *N. reticularis*, *Sowerby*, (*reticulata*, *Benson*) Conch.
 Illustr. f. 44. Calcutta.
 52. *N. obtusa*, *Benson*, Conch. Illustr. f. 43. Calcutta.
 53. *N. Coromandeliana*, *Sowerby*, Conch. Illustr. f. 52.
 Coromandel. (*Humphrey*, ex mus. Portl.)
 54. *N. Philippinarum*, *Sowerby*, Conch. Illustr. f. 53.
 Manilla.
 55. *N. Cassiculum*, *Sowerby*, Conch. Illustr. f. 55.
 56. *N. Sumatrensis*, *Sowerby*, Conch. Illustr. f. 54.
 57. *N. pulchra*, *Sowerby*, Conch. Illustr. f. 59.
 58. *N. triserialis*, *Sowerby*, Conch. Illustr. f. 60.
 59. *N. Guyaquilensis*, *D'Orb.* Guyaquil.

Peritina.

CORRECTED LIST OF FIGURES.

- Fig. 1. *N. picta*, Sow. Sp. 35.
2. *N. reticulata*, Sow. Sp. 36.
3. *N. latissima*, Brod. Sp. 29.
4. *N. Strigilata*, Lam. Sp. 18.
5. *N. cariosa*, Gray, Sp. 26.
6. *N. granosa*, Sow. Sp. 24.
7. *N. intermedia*, Sow. Sp. 33.
8. *N. brevispinosa*, Lam. Sp. 11.
9. *N. spinosa*, Budgin, Sp. 10.
10. *N. Faba*, Sow. Sp. 38.
11. *N. dilatata*, Brod. Sp. 30.
12. *N. globosa*, Brod. Sp. 31.
13. *N. afra*, Sow. Sp. 39.
14. *N. subgranosa*, Sow. Sp. 41.
15. *N. Oweniana*, Gray, Sp. 25.
16. *N. latissima*, Brod. Sp. 29.
17. *N. auriculata*, Lam. Sp. 13.
18. *N. piperina*, Chemn. Sp. 40.
19. *N. Meleagris*, Lam. Sp. 19.
20. *N. Corona*, Lam. Sp. 9.
21. *N. punctulata*, Lam. Sp. 6.
22. *N. canalis*, Sow. Sp. 2.
23. *N. semiconica*, Lam. Sp. 17.
24. *N. viridis*, Lam. Sp. 22.
25. *N. crepidularia*, Lam. Sp. 12.
26. *N. pulligera*, Lam. Sp. 1.
27. *N. virginea*, Lam. Sp. 20.

- Fig. 28. *N. dubia*, *Lam.* Sp. 3.
 29. *N. gagates*, *Lam.* Sp. 7.
 30. *N. pupa*, *Dillwyn*, Sp. 45.
 31. *N. Zebra*, *Lam.* Sp. 4.
 32. *N. strigilata*, *Lam.* var. Sp. 18.
 33. *N. fluviatilis*, *Lam.* Sp. 21.
 34. *N. chlorostoma*, *Sow.* Sp. 34.
 35. *N. fasciata*, *Lam.* Sp. 15.
 36. *N. Smithii*, *Gray*, Sp. 28.
 37. *N. lineolata*, *Lam.* Sp. 16.
 38. *N. lugubris*, *Lam.* Sp. 8.
 39. *N. Nux*, *Brod.* Sp. 32.
 40. *N. Morio*, *Sow.* Sp. 37.
 41. *N. ziczac*, *Lam.* Sp. 5.
 42. *N. Domingensis*, *Lam.* Sp. 14.
 43. *N. obtusa*, *Benson*, Sp. 52.
 44. *N. reticularis*, *Sow.* Sp. 51.
 45. *N. Bætica*, *Lam.* Sp. 23.
 46. *N. Prevostiana*, Sp. 42.
 47. *N. Danubialis*, *Sadler*, Sp. 43.
 48. *N. labiosa*, *Sow.* Sp. 49.
 49. *N. Jordani*, *Sow.* Sp. 48.
 50. *N. subsulcata*, *Sow.* Sp. 50.
 51. *N. Caffra*, *Gray*, Sp. 27.
 52. *N. Coromandeliana*, *Sow.* Sp. 53.
 53. *N. Philippinarum*, *Sow.* Sp. 54.
 54. *N. Sumatrensis*, *Sow.* Sp. 56.
 55. *N. Cassiculum*, *Sow.* Sp. 55.
 56. *N. lutesens*, Sp. 46.
 57. *N. Dalmatica*, *Partch*, Sp. 44.
 58. *N. rivalis*, *Parreyss*, Sp. 47.
 59. *N. pulchra*, *Sow.* Sp. 57.
 60. *N. triserialis*, *Sow.* Sp. 58.

Neritina



1



2



2



3



4



4



3



5




5



1


1. *Alouatta palliata*, Linn. Sp. 1.
 2. *Alouatta palliata*, Linn. Sp. 1.
 3. *Alouatta palliata*, Linn. Sp. 1.
 4. *Alouatta palliata*, Linn. Sp. 1.
 5. *Alouatta palliata*, Linn. Sp. 1.
 6. *Alouatta palliata*, Linn. Sp. 1.
 7. *Alouatta palliata*, Linn. Sp. 1.
 8. *Alouatta palliata*, Linn. Sp. 1.
 9. *Alouatta palliata*, Linn. Sp. 1.
 10. *Alouatta palliata*, Linn. Sp. 1.



Neritina



G.B.S. pin^r exc.



Neritina.

10



11



11



12



9



9



12



13



13



10


6.B.5.jun. exc.



Neritina.



G.B.S. jun. exc.



Neritina



19



21



21



20



22



22



20



23




23




19





Neritina



24



25



25



26



27



27



26



27



27




27




G.B. S. Jun. exc.

Neritina





Neritina



Neritina.







49




49



45



51



50



50



48



47




46



47

unlabeled







A CATALOGUE

OF THE RECENT SPECIES OF THE GENUS

EULIMA.

§ 1. perforatæ.

- Sp. 1 *splendidula nobis*. Conch. Ill. f. 7. St. Elena. Mr. Cuming.
2 *marmorata nobis*. Conch. Ill. f. 8.
3 *interrupta nobis*. Conch. Ill. f. 11. Gulf of Nocoliyo, Mr. Cuming.
4 *imbricata, nobis*. Conch. Ill. f. 4. St. Elena, Mr. Cuming.
5 *brunnea nobis*. Conch. Ill. f. 9. Haynam and China.

§ 2. imperforatæ.

- 6 *brevis nobis*. Conch. Ill. f. 15. Lord Hoods Island, Mr. Cuming.
7 *hastata nobis*, Conch. Ill. f. 10. St. Elena, Mr. Cuming.
8 *major nobis*. Conch. Ill. f. 1. var. f. 1* and f. 1.** Tahiti, Mr. Cuming.
9 *labiosa, nobis*. Conch. Ill. f. 2.
10 *anglica nobis*. Conch. Ill. f. 5. Turbo politus. Mont. British.
11 *subangulata nobis*. Conch. Ill. f. 8. East Indies.
12 *pusilla nobis*. Conch. Ill. f. 6. St. Elena, Mr. Cuming.
13 *articulata nobis*. Conch. Illust. f. 12. Australia.
14 *varians nobis*. Conch. Ill. f. 14. Xipixapi, Mr. Cuming.
15 *lineata nobis*. Conch. Ill. f. 13. British? Turbo glaber Dacosta.

EULIMA.

Corrected list of figures in parts 52, 53, of
CONCHOLOGICAL ILLUSTRATIONS.

Fig. 1. *E. major*.

1.* ——— var.

1** ——— var.

2. *E. labiosa*.

3. *E. subangulata*.

4. *E. imbricata*.

5. *E. anglica*.

6. *E. pusilla*.

7. *E. splendidula*.

8. *E. marmorata*.

9. *E. brunnea*.

10. *E. hastata*.

11. *E. interrupta*.

12. *E. articulata*.

13. *E. lineata*.

14. *E. varians*.

15. *E. brevis*.

Eulina.

2



6



4



1



6



1**



3



5



1*

G.B.S. jun. ex.

Index

1912

Journal of the Royal Asiatic Society

CHRONOLOGICAL ILLUSTRATION

Fig. 1. E. 1000

1. 1000-1000

2. 1000-1000

3. 1000-1000

4. 1000-1000

5. 1000-1000

6. 1000-1000

7. 1000-1000

8. 1000-1000

9. 1000-1000

10. 1000-1000

11. 1000-1000

12. 1000-1000

13. 1000-1000

14. 1000-1000

15. 1000-1000

16. 1000-1000

17. 1000-1000

18. 1000-1000

19. 1000-1000

20. 1000-1000

Eulina.

15



11



9



7



8



12



13



14



10



14

1

2

3

4

5

6

7

8

9

10

A CATALOGUE

OF ALL THE SPECIES HITHERTO KNOWN OF LEACH'S
GENUS MARGARITA,

Represented in the 132nd, 133rd, and 134th Parts of

Conchological Illustrations,

BY

G. B. SOWERBY, JUN.

- Spec. 1. *Margarita tæniata*, Conch. Ill. fig. 2.
2. ——— *violacea*, *King*, in Zool. Journal, vol. v. p. 346.
Conch. Ill. f. 11.
——— *violacea* var. Conch. Ill. f. 12.
3. ——— *expansa*, *Sowerby*, in Malacol. and Conchol.
Mag. p. 24. Conch. Ill. f. 16.
——— *expansa* var. Conch. Ill. f. 17.
4. ——— *sigaretina*, *Sowerby*, in Malacol. and Conchol.
Mag. p. 24. Conch. Ill. f. 14.
5. ——— *vulgaris*, *Leach*, Conch. Ill. f. 13.
6. ——— *carnea*, (*Turbo carneus* *Lowe*) Conch. Ill. f. 9.
7. ——— *striata*, *Brod. & Sowerby*, in Zool. Journal, iv.
Conch. Ill. f. 3. 18.
8. ——— *arctica*, *Leach*, Conch. Ill. f. 6.
9. ——— *grænlantica*, *Beck*, Conch. Ill. f. 10.
10. ——— *umbilicalis*, *Brod. & Sowerby*, Zool. Journal, iv.
Conch. Ill. f. 5.
11. ——— *sulcata*, *Sowerby*, in Malacol. and Conchol.
Mag. p. 26. Conch. Ill. f. 1.
12. ——— *undulata*, *Sowerby*, in Malacol. and Conchol.
Mag. p. 26. Conch. Ill. f. 4.
13. ——— *costellata*, *Sowerby*, in Malacol. and Conchol.
Mag. p. 26. Conch. Ill. f. 15.
14. ——— *acuminata*, *Sowerby*, in Malacol. and Conchol.
Mag. p. 26. Conch. Ill. f. 7.
15. ——— *Solariiformis*, *Sowerby*, in Malacol. and Con-
chol. Mag. p. 26. Conch. Ill. f. 8.

A CATALOGUE

OF ALL THE BOOKS WITHED TO KNOW OR LEARN
CENTRE MARGARITA.

Reprinted in No. 1234, 1235, and 1236.

Contingencies of the

C. R. NORTH, 1894.

1. The first of the series, No. 1, is a...
2. The second of the series, No. 2, is a...
3. The third of the series, No. 3, is a...
4. The fourth of the series, No. 4, is a...
5. The fifth of the series, No. 5, is a...
6. The sixth of the series, No. 6, is a...
7. The seventh of the series, No. 7, is a...
8. The eighth of the series, No. 8, is a...
9. The ninth of the series, No. 9, is a...
10. The tenth of the series, No. 10, is a...
11. The eleventh of the series, No. 11, is a...
12. The twelfth of the series, No. 12, is a...
13. The thirteenth of the series, No. 13, is a...
14. The fourteenth of the series, No. 14, is a...
15. The fifteenth of the series, No. 15, is a...
16. The sixteenth of the series, No. 16, is a...
17. The seventeenth of the series, No. 17, is a...
18. The eighteenth of the series, No. 18, is a...
19. The nineteenth of the series, No. 19, is a...
20. The twentieth of the series, No. 20, is a...

Margarita

G.B.S. fars. ecc

Margarita



9



10



10



5



8



8



5




7



7



6



G. B. 1. 1. 1.

Margarita



12



11



11



13



14



14



14



15



15



16



16



17

4/18/18

A CATALOGUE
OF
RECENT SPECIES OF THE GENUS
CANCELLARIA,

BY

G. B. SOWERBY.

- 1 *Canc. reticulata* Lam. Enycl. meth t. 375 f. 3. Antigua.
- 2 *Canc. candida nobis*, Conch. Illustr. f. 1. Shell ovate oblong, with an acuminate apex, white; volutions seven, entirely covered with decussating ribs; aperture oblong, canal distinct, folds on the columella two, of which the upper is much the larger; edge of the outer lip sharp, inside with distant rib-like teeth; length 1.2, breadth 0.7, Polynesia, Mr. S. Stuchbury.
- 3 *Canc. ovata nobis*, Conch. Illustr. f. 2. Z. P. 1832 p. 53. Dredged in sand at from eight to ten fathoms at St. Elena, Western Columbia, Mr. Cuming.
- 4 *Canc. obesa nobis*, Conch. Illustr. f. 3 jun. f. 4. adult Z. P. 1832, p. 52. From Puerto Portrero and the Gulf of Dulce in central America: dredged at the latter place from a depth of 15 fathoms, Mr. Cuming.
- 5 *Canc. acuminata nobis*, Conch. Illustr. f. 5. Z. P. 1832 p. 53. Dredged in sandy mud at a depth of twelve fathoms at Guacomayo in central America, Mr. Cuming.
- 6 *Canc. solida nobis*, Conch. Illustr. f. 6. Z. P. 1832 p. 50. Found at Real Llejos and St. Elena, by dredging in a sandy bottom at from eight to ten fathoms, Mr. Cuming.
- 7 *Canc. gemmulata, nobis*, Conch. Illustr. f. 7. Z. P. 1832 p. 55. Dredged from a sandy muddy bottom in the Gulf of Nocoioy, in central America, Mr. Cuming.

- 8 *Canc. decussata nobis*, Conch. Illustr. f. 8. Z. P. 1832, p. 55. Found in sandy mud by dredging in 10 to 13 fathoms, at Panama and Puerto Portrero.
- 9 *Canc. indentata, nobis*, Conch. Illustr. f. 9. test. jun. f. 10. adult. Z. P. 1832, p. 54. Panama.
- 10 *Canc. buccinoides nobis*, Conch. Illustr. f. 11. Z. P. 1832, p. 54. Found in sandy mud from 7 to 15 fathoms at Real Llejos, Iquiqui, Callao and Puerto Portrero, by Mr. Cuming.
- 11 *Canc. clavatula nobis*, Conch. Illustr. f. 12. Z. P. 1832, p. 52. Found in sandy mud, in 7 fathoms at Panama and at Payta, Mr. Cuming.
- 12 *Canc. uniplicata nobis*, Conch. Illustr. f. 13. Z. P. 1832. Two specimens were dredged in sand at a depth of ten fathoms, near Panama, by Mr. Cuming.
- 13 *Canc. mitriformis nobis*, Conch. Illustr. f. 14. Z. P. 1832 p. 51. Sandy mud at Panama. Mr. Cuming.
- 14 *Canc. Tritonis nobis*, Conch. Illustr. f. 15. Shell oblong, light brown, varied with darker; with an acuminate spire; volutions six, rather ventricose, longitudinally ribbed and spirally striated, with a slightly nodular-keel near the upper or posterior part, above which it is blotched with dark brown; aperture large and wide; outer lip grooved within; columella rugulose near the anterior part; with three small plaits, of which the middle one is smallest, length 2.3, breadth 1.3. Locality unknown.
- 15 *Canc. granosa nobis*, Conch. Illustr. f. 16, 17. Shell oblong, reddish brown, with an acuminate spire half the length of the shell; volutions six, flattened on the upper part, longitudinally ribbed, and transversely grooved, ribs granose, more strongly so above: aperture wide, outer lip grooved within; columella slightly rugulose in front, with three plaits, the two lower

smaller: canal distinct: length 1.7, breadth 1.0.

Locality unknown.

- 16 *Canc. piscatoria nobis*, *Canc. nodulosa* Lam. Bucc. piscatorium. Chemnitz iv. f. 1151, 1152.
- 17 *Canc. Littoriniformis nobis*, Conch. Illustr. f. 18. Shell subglobose, chesnut brown, rough; spire acuminate; volutions five, rounded with close-set cross striæ; aperture nearly oval, outer lip grooved within, canal very short and indistinct; plaits on the columella three, small; umbilicus rather large, with a rounded edge: length 0.75, breadth 0.5. From Ceylon.
- 18 *Canc. elegans nobis*, Genera of Shells, No. v. fig. 3.
- 19 *Canc. asperella* Lam. Encycl. meth. t. 374 f. 3.
- 20 *Canc. oblonga nobis*, Tank. Catal. app. p. xv. Conch. Illustr. f. 19.
- 21 *Canc. tessellata nobis*, Conch. Illustr. f. 20 jun. f. 20* adult. Z. P. 1832 p. 51. Dredged in sandy mud at a depth of from 7 to 10 fathoms in the Bay of Caraccas, at St. Elena and at Xipixapi. Mr. Cuming.
- 22 *Canc. nodulifera nobis*, Tank. Cat. app. p. xv. No. 1544. Conch. Illustr. f. 21.
- 23 *Canc. Cassidiformus nobis*, Conch. Illustr. f. 22. Z. P. 1832 p. 53. Young shells of this species were dredged in sandy mud in 16 fathoms water, at Panama, by Mr. Cuming.
- 24 *Canc. australis nobis*, Conch. Illustr. f. 23. Shell ovato-oblong, decussated, white; spire rather acuminate, obtuse at the apex; volutions five, rounded with a deep suture, and covered with decussating ribs; aperture oval, outer lip grooved within; columellar lip slightly rugulose, with 3 plaits; canal distinct; length 0.8, breadth 0.5. New South Wales; the late G. Humphrey's Collection.
- 25 *Canc. lævigata nobis*, Conch. Illustr. f. 24. Shell ovato-oblong. smooth, white; spire rather acuminate, obtuse

at its apex ; volutions five, rounded, somewhat flattened at the upper part and with a deep suture ; aperture oval, outer lip deeply grooved within ; columellar lip slightly rugulose in front, with 3 plaits ; canal distinct ; length 0.9, breadth 0.55. New South Wales, the late G. Humphrey.

26 *Canc. spirata* Lam. Conch. Illustr. f. 25. I have received several specimens of this species from Garden Island. Mr. Gray has informed me that the specimens under this name in Lamarck's own collection are not *Cancellariæ* ; the description given in his Hist. nat. des animaux sans vertèbres, accords, however, so perfectly with my shells, that I cannot hesitate to regard them as Lamarck's species.

27 *Canc. obliquata* Lam. Conch. Illustr. f. 26. Haynam.

28 *Canc. scalata nobis*, Conch. Illustr. f. 27. Shell oblong, reddish brown, spire acuminate ; volutions six, with a bluntly angular-edged and broad groove next to the suture, the margin of which groove is crenulate ; the volutions covered with longitudinal, somewhat nodulose ribs, the interstices of which have a few cross-striæ ; aperture white, oval, truncated and with an internal tubercle at the upper part, outer lip grooved within ; columella with four small plaits, rugulose, umbilicus small, narrow ; length 1.2, breadth 0.7. East Indies.

29 *Canc. contabulata nobis*. Conch. Illustr. f. 28. Shell oval, pale brown, nodulose-ribbed ; spire rather short ; volutions five, ventricose, their upper part next the suture flat, with a bluntly angular edge ; nodulose longitudinal ribs continued obliquely over the flat space to the suture ; aperture large, white, oval, truncated at its upper part ; outer lip grooved within ; columella with four small plaits, of which the upper one is very small ; umbilicus small : length 1.2, breadth 0.8. Ceylon.

- 30 *Canc. crenifera nobis*, Conch. Illustr. f. 29. Shell ovate; pinkish brown, shining; volutions 5 to 6, rounded with longitudinal crenulated ribs, the upper ends of which form a strongly crenated edge to the canaliculated suture; aperture oval, white; outer lip thickened, reflected, grooved within, columella with three plaits; canal indistinct; umbilicus small but distinct and rough at its edge: length 0.9, breadth 0.5. From Ceylon.
- 31 *Canc. scalarina*. Lam. Chemn. iv. f. 1172, 1173.
- 32 *Canc. crista nobis*, Conch. Illustr. f. 30. Shell oval, brownish with lighter bands, rough; spire acuminate; volutions 5 to 6, rather ventricose, with longitudinal ribs, which are dentated on their edges and raised above the edge of the canaliculated suture, aperture ovate, truncated at the upper part and forming a small canal anteriorly; outer lip grooved within; plaits on the columella three; umbilicus small: length 1.0, breadth 0.6. Locality unknown.
- 33 *Canc. costifera, nobis*, Conch. Illustr. f. 31. Shell oval, light brown, spire acuminate, volutions six, angular at the upper edge, flat next to the suture, with distant longitudinal ribs, which are pointed at the upper part, and then continuous in oblique direction across the flat upper edge of the volutions: interstices slightly cross-striated: aperture oval, truncated at the upper part, outer lip grooved within; columella with three plaits, umbilicus very small: length 0.8, breadth, 0.45.
- 34 *Canc. articularis nobis*, Conch. Illustr. f. 32. Shell oblong, subturrit, orange brown; spire acuminate; volutions six, canaliculated at the suture, obliquely longitudinally ribbed, ribs prominent at the edge of the channelled suture and crenated on their edges; aperture wide, oval, truncated at the upper part; outer lip reflected, finely grooved within; columella with three small plaits; umbilicus small; length 0.9, breadth 0.5. Ceylon.

- 35 *Canc. brevis nobis*, Conch. Illustr. 33. Z. P. 1832 p. 52. Found at Puerto Portrero and St. Elena; Mr. Cuming.
- 36 *Canc. pusilla nobis*, Conch. Illustr. f. 34. Shell subglobose, ventricose, pale; volutions four, with a broad flat area at the upper part, which is crenated at both edges; longitudinally ribbed and cross-straited; aperture trigonal, plaits on the columella two; umbilicus none: length 0.5, breadth 0.4, a single specimen was taken out of the umbilicus of a *Cassis Rufa* from New South Wales.
- 37 *Canc. bullata nobis*, Conch. Illustr. f. 35. Z. P. 1832 p. 51. Dredged in mud at a depth of twelve fathoms, at Payta, and in the Gulf of Nocoliyo. Mr. Cuming.
- 38 *Canc. tuberculosa nobis*, Conch. Illustr. f. 36. Z. P. 1832 p. 51. Dredged in sandy mud, at a depth of seven fathoms, at Iquique. Mr. Cuming.
- 39 *Canc. pulchra nobis*, Conch. Illustr. f. 37. Z. P. 1832 p. 50. Dredged from a sandy bottom in from eight to ten fathoms, at St. Elena. Mr. Cuming.
- 40 *Canc. cancellata* Lam. Encycl. Method. 375. f. 4.
- 41 *Canc. similis nobis*, Conch. Illustr. f. 38. Shell ovate, rather compressed, varicose; spire somewhat acuminate, volutions six to seven, gradually increasing, whitish with irregular brown bands, longitudinally obliquely ribbed, ribs distant, crossed by strongly marked elevated lines; aperture white, ringent, outer lip varicose outside, crenated at its border, deeply grooved and strongly toothed a little way within the edge, columella with three plaits, of which the lowest is the smallest and double; canal reflected, umbilicus very small with a broad and distinct border; length 1.2, breadth 0.9.
- 42 *Canc. chrysostoma nobis*, Conch. Illustr. f. 39. Z. P. 1832, p. 54. Found in sand at from eight to ten fathoms, at Panama and St. Elena. Mr. Cuming.

- 43 *Canc. rugosa* Lam. Encycl. meth. t. 375 f. 8. China.
- 44 *Canc. hæmastoma nobis*, Conch. Illustr. f. 40. Z. P. 1832, p. 51. Found in sand at from 10 to 16 fathoms, near the Galapagos Islands. Mr. Cuming.
- 45 *Canc. rigida nobis*, Conch. Illustr. f. 41. Z. P. 1832, p. 53. Dredged in sand at 13 fathoms deep at Puerto Portrero. Mr. Cuming.
- 46 *Canc. costata* Gray, Conch. Illustr. f. 42. Shell oblong, fulvous varied and banded with dark brown, longitudinally ribbed, and transversely striated; spire acuminate, volutions six, with a flat, sharp-edged space at the upper or posterior part; ribs distant, sharp-edged anteriorly, continuing obliquely over the flat space to the suture; aperture subtrigonal, outer lip slightly grooved within; columella with two distinct and one very small plait, lip a little rugose; umbilicus rather large; length 0.95, breadth 0.7.
- 47 *Canc. goniosstoma nobis*, Conch. Illustr. f. 43. Chemn. iv. Vign. 37 f. a. b. c. Z. P. 1832, p. 51. A single specimen was brought up from a sandy bottom in eight fathoms water at Conchagua by Mr. Cuming.
- 48 *Canc. trigonostoma nobis*, Conch. Illustr. f. 44. *Delphinula trigonostoma* Lam. vi. pt. ii. p. 231. Shell pyramidal, acuminate, white or pale brown; volutions five, flat and spirally striated on the upper part; the edge of which is strongly keeled and crenated; slightly longitudinally ribbed and spirally striated on the lower part; aperture triangular, with a distinct varicose edge, and three small plaits on the columella; umbilicus very large, crenated at the edge; length 0.9, breadth 0.75. Ceylon.

Obs. It is remarkable that in this shell, which is the *Bordstrappe* of the Dutch Conchologists, the inner edges of the volutions are set against the edge of the umbilicus.

CANCELLARIÆ.

Corrected List of Figures, in parts 9 to 13.

- | | |
|--------------------------------|------------------------------|
| Fig. 1. <i>C. candida.</i> | Fig. 24. <i>C. lævigata.</i> |
| 2. <i>C. ovata.</i> | 25. <i>C. spirata.</i> |
| 3 & 4. <i>C. obesa.</i> | 26. <i>C. obliquata.</i> |
| 5. <i>C. acuminata.</i> | 27. <i>C. scalata.</i> |
| 6. <i>C. solida.</i> | 28. <i>C. contabulata.</i> |
| 7. <i>C. gemmulata.</i> | 29. <i>C. crenifera.</i> |
| 8. <i>C. decussata.</i> | 30. <i>C. crispa.</i> |
| 9. <i>C. indentata, jun.</i> | 31. <i>C. costifera.</i> |
| 10. <i>C. indentata.</i> | 32. <i>C. articularis.</i> |
| 11. <i>C. buccinoides.</i> | 33. <i>C. brevis.</i> |
| 12. <i>C. clavatula.</i> | 34. <i>C. pusilla.</i> |
| 13. <i>C. uniplicata.</i> | 35. <i>C. bullata.</i> |
| 14. <i>C. mitriformis.</i> | 36. <i>C. tuberculosa.</i> |
| 15. <i>C. Tritonis.</i> | 37. <i>C. pulchra.</i> |
| 16, 17. <i>C. granosa.</i> | 38. <i>C. similis.</i> |
| 18. <i>C. Littoriniformis.</i> | 39. <i>C. chrysostoma.</i> |
| 19. <i>C. oblonga.</i> | 40. <i>C. hæmastoma.</i> |
| 20, 20.* <i>tessellata.</i> | 41. <i>C. rigida.</i> |
| 21. <i>C. nodulifera.</i> | 42. <i>C. costata.</i> |
| 22. <i>C. cassidiformis.</i> | 43. <i>C. goniostoma.</i> |
| 23. <i>C. Australis.</i> | 44. <i>C. trigonostoma.</i> |

Obs. Lamarck's *C. senticosa*, *C. citharella*, and *C. Ziervogliana* do not belong to this genus; the first is a *Phos*, the 2nd, an *Oniscia*, and the last a *Mitra*. His *C. scalariformis* is unknown to the author.

Cancellaria.

Cancellaria.





Cancellaria.





Cancellaria.

34



29



29



33



28



35



32



30




31




33



Cancellaria.





MUREX.

A CATALOGUE OF RECENT SPECIES,

BY

G. B. SOWERBY, Jun.

Sp.

1. *M. tenuispina*, *Lam.* vii. p. 158. *Ch. Conch.* t. 189. f. 1821. t. 190. f. 1822. Moluccas. *Gray* says that the *M. tenuirostrum* of *Lam.* is a specimen of this, without spines. Philippines.
2. *M. ternispina* *Lam.* vii. p. 158. *Con. Illus.* f. 110 *Var.* *M. aduncospinosus*, *Beck.* *Con. Illus.* f. 68. E. Indies.
3. *M. crassispina*, *Lam.* vii. p. 157. *Martini.* t. 113. f. 1052. Philippines. In this species the 2nd spine is longest.
4. *M. rarispina*, *Lam.* 7. p. 158, *M. unidentatus*, *Con. Illus.* f. 52, E. Indies.
5. *M. Scolopax*, *Dillwyn*, *Descr. Cat.* p. 681. *Ch. Conch.* f. 1819—20. Red Sea. Smoother than the last, with yellowish brown lines across.
6. *M. Occa* *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 45. Nicobar Isl.
7. *M. brevispina*, *Lam.* vii. p. 159. *Con. Illus.* f. 10.
8. *M. chrysostoma*, *Gray?* *Con. Illus.* f. 1.
9. *M. Messorius*. *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 93.
10. *M. Haustellum*, *Lim. Gmel.* p. 3524. *Martini.* *Conch.* t. 115. f. 1066. Moluccas. Philippines.
11. *M. rectirostrum*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 111.
12. *M. nigrescens*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 113. Xipixapi. Columbia. Mr. Cuming.
13. *M. recurvirostrum*, *Brod. Zool. Proc.* 1832. p. 174. *Con. Illus.* f. 9. Gulf of Nocoyo. Central Amer. Mr. Cuming.
14. *M. pliciferus*. *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 101.

- Sp.
 15. *M. plicatus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 6.
 Gulf of Nocoyo. Mr. Cuming.
16. *M. formosus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus.
 f. 112. Persian Gulf.
17. *M. Mindinaoensis*. *Sow. Jun. Zool. Proc.* 1840. Con.
 Illus. f. 92. Misamis, Mindinao, Philippines. Mr.
 Cuming.
18. *M. Motacilla*, *Linn. Gmel.* p. 3530. Chemn. Conch. t.
 163. f. 1563. Indian Ocean. *Var.* Con. Illus. f. 69.
19. *M. elegans*, *Beck. Sow. Jun. Zool. Proc.* 1840. Con.
 Illus. f. 84.
20. *M. similis*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 70.
21. *M. inflatus*, *Lam. vii.* p. 160. Martini. t. 102. f. 980. t.
 103. f. 981. E. Indies. Has a triangular tooth on
 the outer lip.
22. *M. elongatus*. *Lam. vii.* p. 161. Con. Illus. f. 88. *Var.* Con.
 Illus. f. 89. E. Indies. Has the tooth as in *M. inflatus*.
23. *M. anguliferus*, *Lam. vii.* p. 171. Martini. t. 110. f.
 1029—1030. *M. inflatus*, *Var.* Con. Illus. f. 53. *M.*
Ferrugo, *Wood. Index, Test. costatus?* *Gmel.* Not
 including the "Sirat" of Adanson. Atlantic Ocean.
24. *M. Senegalensis*, *Gmel.* "Sirat" *Adanson.* *M. Brasili-*
ensis, Con. Illus. f. 55, white variety, Con. Illus.
 f. 61. Senegal.
25. *M. Calcitraba*, *Lam. vii.* p. 161. Martini. t. 103. f. 982
M. brevifrons. *Lam. vii.* 160. Martini, t. 103. f.
 983. t. 104, f. 984—6. America. Subject to great
 variations, but distinguished from *M. inflatus* by
 the absence of the tooth. Red Sea, Indian Ocean.
26. *M. Monodon.* *Sow. Tank. Cat.* Martini, iii. t. 105, f.
 987-8. Swan River.
27. *M. Pomum*, *Gmel.* p. 3327. *M. asperrimus*, *Lam. vii.* p.
 164. Martini; t. 109. f. 1021-1023. West Indies.
28. *M. scabrosus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 73.
29. *M. Cervicornis*, *Lam. vii.* p. 163, *Sow. Gen. Murex.* f. 2.
 New Holland.

- Sp.
 30. *M. axicornis*, *Lam.* vii. p. 163. Martini Conch. t. 105.
 f. 989. var. black, Con. Illus. f. 66. Moluccas.
 Philippines.
31. *M. Banksii*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 82.
32. *M. aculeatus*, *Lam.* Con. Illus. f. 63. Moluccas.
33. *M. Palma-rosæ*, *Lam.* vii. p. 161. Lister, Conch. t. 946.
 f. 41. Ceylon.
34. *M. Saulii*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 77.
 A dwarf variety is brought from Ticao by Mr.
 Cuming.
35. *M. torrefactus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus.
 120; *Var.* 121. Ticao, Philippines. Mr. Cuming.
36. *M. adustus*, *Lam.* vii. p. 162. Martini. t. 105, f. 990
 991. Singapore and Philippines.
37. *M. rubescens*, *Brod. Zool. Proc.* 1832. p. 174. Con. Illus.
 f. 7 Tahiti. Mr. Cuming.
38. *M. microphyllus*, *Lam.* vii. p. 163. E. M. pl. 415. f. 5.
 Con. Illus. f. 105.
39. *M. Maurus*, *Brod. Zool. Proc.* 1832. p. 174. Con. Illus. f.
 12. *Var.* f. 67. Isl. Annaa, Pacific Ocean.
40. *M. Capucinus*, *Lam.* vii. p. 164. Chem. Conch. t. 192.
 f. 1849. 1850. Nicobar Isl. E. Indies. Singapore.
41. *M. castaneus*, Con. Illus. f. 44. Mr. Norris's collection.
42. *M. rufus*, *Lam.* vii. p. 162. White variety, f. 99. Indian Seas.
43. *M. palmiferus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus.
 f. 104. Red Sea.
44. *M. corrugatus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus.
 f. 72.
45. *M. laqueatus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 78.
46. *M. triqueter*, *Born.* p. 291. t. 11. f. 1-2. Martini, t. 111.
 f. 1038. Philippines.
47. *M. laciniatus*, *Nob. Zool. Proc.* 1841. Con. Illus. f. 59.
48. *M. trigonulus*, *Lam.* vii. p. 167. Con. Illus. f. 102.
 Persian Gulph.
49. *M. pinniger*, *Brod. Zool. Proc.* 1832. p. 174. Con. Illus.
 f. 50 and 109. (*M. cristatus*) Xipixapi. Mr. Cuming.
50. *M. cancellatus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 79.

- Sp.
 51. *M. canaliferus*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 74.
52. *M. Mitriformis*, *Sow. Jun. Zool. Proc.* 1841. *Con. Illus.* f. 75.
53. *M. Capensis*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 76. Cape of Good Hope.
54. *M. pinnatus*, *Swainson, Bligh. Cat. Appendix* p. 17. *Zool. Illus.* 2nd Ser : pl. 122. China.
55. *M. tripterus*, *Born*, p. 291. t. 10. f. 18-19. *Wood, Index. Test. M. Trialatus*, *Con. Illus.* f. 54. 54* (Not *M. Trialatus*, *C. I.* f. 33.) Gmelin was mistaken in referring this sp. to the fossil which bears the name. Ticao, Philippines.
56. *M. trigonularis*, *Lam. vii.* p. 165. *Con. Illus.* f. 107. Our shell agrees so well with Lamarck's description, that we venture to apply the name to it, notwithstanding Mr. Gray's statement, that the *M. trigonularis* of *Lam.* is nothing but *Acanthopterus* deprived of its fringes. Mr. Gray cannot have studied the subject, for he has mixed up the latter species with the "*M. Purpura alata*" of *Chemnitz*. (*M. foliatus* of *Gmelin*.) Bantayan, Philippines. Mr. Cuming.
57. *M. uncinarius*, *Lam. vii.* p. 166. *Martini iii.* f. 1034-5. *Con. Illus.* f. 106. Isl. Masbate, Philippines. Mr. Cuming.
58. *M. trialatus*, *Sow. Jun. Zool. Proc.* 1841. *Con. Illus.* f. 33. (Not *trialatus*, f. 54. 54.*)
59. *M. acanthopterus*, *Lam. vii.* f. 165. *E. M.* pl. 417. f. 2. a. b. *Con. Illus.* f. 85. Brown variety, young (*alatus*), *Con. Illus.* f. 51. A very large shell from Isl. Mindinao exists in Mr. Cuming's cabinet, which may probably be referable to this species, *Con. Illus.* f. 115.
60. *M. foliatus* *Gmel.* p. 3529. *M. Tripterus. Lam. vii.* p. 165. "*M. Purpura alata*" *Chem.* 10. t. 161. f. 1538-9 N.W. Co. of America.

- Sp.
 61. *M. emarginatus*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* 98, 100. Mr. Cuming.
 62. *M. monoceros*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus.* f. 64, 65. and 97, *Mus. Norris. Stainforth. Cuming.*
 63. *M. erinaceus*, *Lam. vii. p. 172. Martini. t. 110. f. 1026-8.* *M. decussatus*, *Gmel. Britain. Mediterranean.*
 64. *M. Tarentinus*, *Lam. vii. p. 173. M. erinaceus*, var. *Con. Illus. f. 24. Sicily, Portugal.* We are still doubtful about this being distinct from *M. erinaceus*.
 65. *M. torosus*, *Lam. vii. p. 174. Con. Illus. f. 39.* Perhaps a variety of the last. Sicily?
 66. *M. Edwardsii*, *Menke. Syn. p. 66. Philippi. Enum. Mollusc. Siciliae. p. 210. Purpura Lassaigii, Basterot. Con. Illus. f. 80. var?* Sicily. Placed here because it has not three regular varices.
 67. *M. inconspicuus*, *Sow. Jun. Zool. Proc.* 1841. *Con. Illus. f. 81. Var. 117. Jersey.*
 68. *M. fasciatus*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus. f. 86, (M. fasciatus, Risso?) R. Gambia.*
 69. *M. varius*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus. f. 57. 108.*
 A variety occurs with more than 3 varices. R. Gambia.
 70. *M. cornutus. Gmel. p. 3525. Martini. Conch. t. 114, f. 1057. Moluccas.*
 71. *M. tumulosus*, *Sow. Jun. Zool. Proc.* 1840. *Con. Illus. f. 71.*
 72. *M. Brandaris. Gmel. p. 3526. Chemn. t. 164. f. 1571. Mediterranean and Adriatic.*
 73. *M. Rota*, — ? *Con. Illus. f. 119. Isle of Capul, Philippines.*
 74. *M. Scorpio. Lam. vii. 169. Martini, t. 106. f. 998-1003. E. Indies. Moluccas, Philippines.*
 75. *M. secundus, Lam. vii. p. 169. Con. Illus. f. 116. Isl. Masbate, Philippines.*
 76. *M. varicosus, Sow. Jun. Zool. Proc.* 1840. *Con. Illus. f. 49.*

- No.
 77. *M. digitatus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 114. Red Sea.
 78. *M. fenestratus*, *Lam.* vii. p. 174. Chemn. 161. f. 1536-1537. Con. Illus. f. 41. Isl. Capul. Philippines.
 79. *M. Jatonus*, *Adanson*. Senegal. 1. t. 9. f. 21. *M. "Lingua vervecina,"* Chem. t. 161. f. 1540. 1. *M. Lingua*, *Dillwyn*, Wood's Index. *M. Gibbosus*, *Lam.* vii. p. 166. In a young state, *M. hemitripterus*, *Lam.* vii. p. 166. E. M. 418. f. 4. a. b. Con. Illus. f. 60. Senegal.
 80. *M. Oxyacantha*, *Brod. Zool. Proc.* 1832. p. 176. Con. Illus. f. 11. Real Llejos, Central America. Mr. Cuming.
 81. *M. humilis*, *Brod. Zool. Proc.* 1832. p. 175. Con. Illus. f. 46, 47. St. Elena. Mr. Cuming.
 82. *M. melanomathos*, *Lam.* vii. p. 168. Martini. 108. p. 1015. E. M. 418. 2, a. b. E. Indies.
 83. *M. Princeps*, *Brod. Zool. Proc.* 1832. p. 175. Con. Illus. f. 43. Puerto Portrero, Central America.
 84. *M. nitidus*, *Brod. Zool. Proc.* 1832. p. 175. Con. Illus. f. 4. Real Llejos, Central America. Probably a variety of the last in a young state.
 85. *M. Radix*. *Gmel.* Swains. Zool. Illus, "Muricinae." Columbia, Bay of Caraccas.
 86. *M. saxatilis*, *Gmel.* p. 3529. Martini 108. f. 1013, 1014. Chemn. t. 161. f. 1528-9. *M. eurystomus*, Sw. Gambia.
 87. *M. Rosarium*. *Chemn.* t. 161. f. 1528-9. Con. Illus. f. 118. *M. Melonulus*, *Lam.* vii. p. 171. E. Indies.
 88. *M. Brassica*, *Lam.* vii, p. 167. *M. Ducalis*, *Brod. Zool. Journ.* iv. p. 377. Con. Illus. f. 56. Mazatlan, Pacific. The description by Lamarck agrees perfectly well with our shell, besides which we have the direct testimony of Mr. Gray, who has seen Lamarck's shell, as to its identity.
 89. *M. Regius*, *Swainson*, Exotic Conch. part 2. Wood's Supplement, Murex, No. 13. Panama.

Sp.

90. *M. Imperialis*, *Swainson* Zool. Illus. "Muricinæ" *M. Pomum*. *Var.* Gmel. p. 3327. Chemn. t. 110, f. 1024-5. California.
91. *M. erythrostroma*, *Swains.* Zool. Illus. "Muricinæ" *Var.* Con. Illus. f. 38. Acapulco.
92. *M. Endivia*, *Lam.* vii. p. 168. *Martini.* t. 107. f. 1008, *M. Saxatilis*, *Wood.* Index. *M. chicoreum*, Gmel? Philippines.
93. *M. Saxicola*, *Brod.* Zool. Journ. Vol. 2. p. 202. Tab. suppl. xi. f. 3. More angulated than *M. Endivia*. Isl. Ticao. Mr. Cuming.
94. *M. turbinatus*, *Lam.* vii. p. 170. Con. Illus. (*M. fulvescens*) f. 30. *Var.* Con. Illus. f. 90, 91.
95. *M. Trunculus*, *Lam.* vii. p. 170. *Martini*, Conch. t. 109. f. 1018, 1020. Mediterranean and Atlantic.
96. *M. angularis*, *Lam.* vii. p. 174. *M. costatus* E. M. 441. f. 3. a b. "Cofar" Adanson. Senegal, 9. f. 22. *M. Octonus*, Con. Illus. f. 32. Senegal.
97. *M. Megacerus*, *Sow. Jun.* Zool. Proc. 1840. Con. Illus. f. 18. S. Seas.
98. *M. Lappa*, *Brod.* Zool. Proc. 1832. p. 177. Con. Illus. f. 15. St. Elena. Mr. Cuming.
99. *M. Zelandicus*, *Quoy et Gaimard*, *Voy. de. l'Astr.* t. 2. p. 529. Pl. 36. f. 5-7. *M. Nova Zelandiæ*, Con. Illus. f. 34. N. Zealand.
100. *M. aduncus*, *Con. Illus.* f. 35 In Mr. Norris's Cabinet.
101. *M. falcatus* *Sow. Jun.* Zool. Proc. 1840. Con. Illus. f. 31. Japan.
102. *M. hexagonus*, *Lam.* vii. p. 169. E. M. pl. 418. f. 3. a. b.
103. *M. inermis*, *Sow. Jun.* Zool. Proc. 1840. Con. Illus. f. 87. Japan.
104. *M. erosus*, *Brod.* Zool. Proc. 1840. Con. Illus. f. 16. var. f. 96. Panama, H. Cuming.
105. *M. balteatus*, *Beck. Sow. Jun.* Zool. Proc. 1840. Con. Illus. f. 83. Masbate, Philippines. H. Cuming.
106. *M. exiguus*, *Brod.* Zool. Proc. 1832. p. 175. Con. Illus. f. 17. Salango. Columbia. Mr. Cuming.

- Sp.
 107. *M. Cyclostoma*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 95. Bohol. Philippines. Mr. Cuming.
 108. *M. lugubris*, *Brod. Zool. Proc.* 1832. p. 175. Con. Illus. f. 26. Puerto Portrero, Central America. Mr. Cuming.
 109. *M. incisus*, *Brod. Zool. Proc.* 1832. p. 176. Con. Illus. f. 13. St. Elena. Mr. Cuming.
 110. *M. breviculus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 37.
 111. *M. tetragonus*, *Brod. Zool. Proc.* 1832. p. 174. Con. Illus. f. 25, 36.
 112. *M. Peruvianus*, *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 103. Pacosmayo, Peru.
 113. *M. Dipsacus*, *Brod. Zool. Proc.* 1832. p. 194. Con. Illus. f. 20. (*M. Octonus*, *Gray. N. Zealand.*) St. Elena.
 114. *M. crispus*, *Brod. Zool. Proc.* 1832. p. 176. Con. Illus. f. 8. *M. tortuosus*. Pacosmayo, Peru.
-

The following are admitted among the Murices with considerable hesitation.

115. *M. miliaris*, *Gmel.* p. 3536. No. 39. *M. vitulinus*, *Lam.* vii. p. 174. *E. M.* pl. 419. f. 1. a. b. 7. a. b.
 116. *M. salebrosus*, *King. Zool. Journ.* v. p. 347. Con. Illus. f. 5. var. f. 48. Southern Coast of S. America.
 117. *M. Fiscellum*, *Chemn.* 10. t. 160. f. 1524. 5. *M. Funiculum*, *Wood's Index. test.* *M. fimbriatus*, *Lam. Iostoma*, Con. Illus. f. 42. *M. Australis*, *Quoy et Gaimard Voy. de l'Astr.* 2. p. 534. pl. 36. f. 13-16. Named *M. ricinuloides* on the plate. *Var.* Con. Illus. f. 62. Pulo Condore.
 118. *M. margariticola*, *Brod. Zool. Proc.* 1832. p. 177. Con. Illus. f. 21. Lord Hood's Isl. Pacific. Mr. Cuming.
 119. *M. dubius*, *Nob. M. aculeatus*, *Wood.* (pre-occupied) *Index. test.* Con. Illus. f. 23. Panama, Mr. Cuming.
 120. *M. noduliferus* *Sow. Jun. Zool. Proc.* 1840. Con. Illus. f. 94. Masbate, Philippines. Mr. Cuming.

121. *M. cristatus*, *Brocchi*, p. 394. t. vii.. f. 15. *Cancellaria cristata*, *Brown*. *Reise* p. 550. *Murex Blainvillii*, *Payradeau*, p. 149. t. 7. f. 17, 18. *M. pliciferus*, *Bivon*, l. c. p. 26. t. 3. f. 10. *Var.* "costatus, nodulis inermibus." *Philippi*, v. t. xi. f. 25. *M. fortis*, *Risso*. *Var.* *M. cataphractus*, *Con.* *Illus.* f. 40. The recent varieties correspond with those in a fossil state, and there is an indivisible chain running from those which are frondose and almost spinose, to those which are characterized by plain simple nodules. Mediterranean.
122. *M. vittatus*, *Broderip*. *Zool. Proc.* 1832. p. 176. *Con.* *Illus.* f. 19. *Isl. Muerte*, Bay of Guayaquil. *Mr. Cuming*.

The following Murices of Authors belong to other Genera, viz:—

FUSI, *M. magellanicus*, *lamellosus*, *scaber*, *lyratus*, and *cingulatus*, of *Lamarck*. *M. Horridus*, *buxeus*, *Carduus squamosus*, *Nucleus*, *cariniferus*, and *pallidus*, of *Broderip*. *M. distinctus*, *Philippi*?

PURPURÆ. *M. costularis*, *Lam.* *M. crispatus*, *Lam.* *M. crassilabrum*, *Gray*.

TYPHIS, *M. fistulosus*, *Brocchi*. *Philippi*.

POLLIE, *M. concatenatus*, *granarius* and *pulchellus*, of *Lamarck*; *M. Vibex*, *Broderip*.

The following cannot be identified.

Lamarck; *M. aciculatus*, *phyllopterus*, *quadrifrons*, *polygonulus*, *cinguliferus* and *subcarinatus*. *King*; *M. Rhodocheilus*, *Z. J. v.* p. 347. *Broderip*; *M. pumilus* *Z. P.* 1832. p. 175. *Conrad*; *M. Nuttali*. *Risso*; *M. fistulatus*, *ramulosus*, *bicristatus*, *Orbignianus*, *rudis*, *bicolor*, *affinis*, *angulatus*, *Pairaudoce*, *triangularis*, *imbricatus*, *postdiluvianus*, *succinctus*, *columnus*. The latter are very vaguely described and neither figures nor synonyms referred to.

MURICES.

Index to the Figures in Parts 58 to 67, and 187 to 199.

(The Figures after the name refers to the Catalogue.)

- | | |
|--|---|
| FIG. | FIG. |
| 1 <i>M. chrysostoma</i> , Sp. 8. | 34 <i>M. Zelandicus</i> , Sp. 99. |
| 2 <i>Fusus nucleus</i> . | 35 <i>M. aduncus</i> , Sp. 100. |
| 3 <i>Fusus pallidus</i> . | 36 <i>M. tetragonus</i> , Sp. 111. |
| 4 <i>M. nitidus</i> , Sp. 84. | 37 <i>M. breviculus</i> , Sp. 110. |
| 5 <i>M. salebrosus</i> , var. Sp. 116. | 38 <i>M. erythrostoma</i> , var. Sp. 91. |
| 6 <i>M. plicatus</i> , Sp. 15. | 39 <i>M. torosus</i> , Sp. 65. |
| 7 <i>M. rubescens</i> , Sp. 37. | 40 <i>M. cristatus</i> , Sp. 121. |
| 8 <i>M. crispus</i> , Sp. 114. | 41 <i>M. fenestratus</i> , Sp. 78. |
| 9 <i>M. recurvirostrum</i> , Sp. 13. | 42 <i>M. Fiscellum</i> , Sp. 117. |
| 10 <i>M. brevispina</i> , Sp. 7. | 43 <i>M. Princeps</i> , Sp. 83. |
| 11 <i>M. Oxyacantha</i> , Sp. 80. | 44 <i>M. castaneus</i> , Sp. 41. |
| 12 <i>M. Maurus</i> , Sp. 39. | 45 <i>M. Occa</i> , Sp. 6. |
| 13 <i>M. insciscus</i> , Sp. 109. | 46 } <i>M. humilis</i> , Sp. 81. |
| 14 <i>Purpura, crassilabrum</i> . | 47 } |
| 15 <i>M. Lappa</i> , Sp. 98. | 48 <i>M. salebrosus</i> , Sp. 116. |
| 16 <i>M. erosus</i> , Sp. 104. | 49 <i>M. varicosus</i> , Sp. 76. |
| 17 <i>M. exiguus</i> , Sp. 106. | 50 <i>M. pinniger</i> , Sp. 49. |
| 18 <i>M. megacerus</i> , Sp. 97. | 51 <i>M. acanthopterus</i> , var. Sp. 59. |
| 19 <i>M. vittatus</i> , Sp. 122. | 52 <i>M. rarispina</i> , Sp. 4. |
| 20 <i>M. dipsacus</i> , Sp. 113. | 53 <i>M. anguliferus</i> , Sp. 23. |
| 21 <i>M. margariticola</i> , Sp. 118. | 54 <i>M. tripterus</i> , Sp. 55. |
| 22 <i>Fusus carduus</i> . | 55 <i>M. Senegalensis</i> , Sp. 24. |
| 23 <i>M. dubius</i> , Sp. 119. | 56 <i>M. Brassica</i> , Sp. 88. |
| 24 <i>M. Tarentinus</i> , Sp. 64. | 57 <i>M. varius</i> , Sp. 69. |
| 25 <i>M. tetragonus</i> , Sp. 111. | 58 <i>Fusus cariniferus</i> . |
| 26 <i>M. lugubris</i> , Sp. 108. | 59 <i>M. laciniatus</i> , Sp. 47. |
| 27 <i>Fusus squamosus</i> . | 60 <i>M. Jatonus</i> , young, Sp. 79. |
| 28 <i>Polia buxea</i> . | 61 <i>M. Senegalensis</i> , var. Sp. 24. |
| 29 <i>Fusus horridus</i> . | 62 <i>M. Fiscellum</i> , var. Sp. 117. |
| 30 <i>M. turbinatus</i> , Sp. 94. | 63 <i>M. aculeatus</i> , Sp. 32. |
| 31 <i>M. falcatus</i> , Sp. 101. | 64 <i>M. Monoceros</i> , Sp. 62. |
| 32 <i>M. angularis</i> , Sp. 96. | 65 front view of the same. |
| 33 <i>M. trialatus</i> , Sp. 58. | 66 <i>M. axicornis</i> var. Sp. 30. |

FIG.

- 67 *M. Maurus*, var. Sp. 39.
- 68 *M. ternispina*, var. Sp. 2.
- 69 *M. Motacilla*, var. Sp. 18.
- 70 *M. similis*, Sp. 20.
- 71 *M. tumulosus*, Sp. 71.
- 72 *M. corrugatus*, Sp. 44.
- 73 *M. scabrosus*, Sp. 28.
- 74 *M. canaliferus*, Sp. 51.
- 75 *M. mitriformis*, Sp. 52.
- 76 *M. Capensis*, Sp. 53.
- 77 *M. Saulii*, Sp. 34.
- 78 *M. laqueatus*, Sp. 45.
- 79 *M. cancellatus*, Sp. 50.
- 80 *M. Edwardsii*, Sp. 66.
- 81 *M. inconspicuus*, Sp. 67.
- 82 *M. Banksii*, Sp. 31.
- 83 *M. Balteatus*, Sp. 105.
- 84 *M. elegans*, Sp. 19.
- 85 *M. acanthopterus*, Sp. 59.
- 86 *M. fasciatus*, Sp. 68.
- 87 *M. inermis*, Sp. 103.
- 88 *M. elongatus*, var. Sp. 22.
- 89 *M. elongatus*, var. Sp. 22.
- 90 *M. turbinatus*, Sp. 94.
- 91 variety of the same.
- 92 *M. Mindinaoensis*, Sp. 17.
- 93 *M. Messorius*, Sp. 9.
- 94 *M. noduliferus*, Sp. 120.

FIG.

- 95 *M. Cyclostoma*, Sp. 107.
- 96 *M. erosus*, var. Sp. 104.
- 97 *M. Monoceros*, Sp. 62.
- 98 *M. emarginatus*, Sp. 61.
- 99 *M. rufus*, var. Sp. 42.
- 100 *M. emarginatus*, Sp. 61.
- 101 *M. pliciferus*, Sp. 14.
- 102 *M. trigonulus*, Sp. 48.
- 103 *M. Peruvianus*, Sp. 112.
- 104 *M. palmiferus*, Sp. 43.
- 105 *M. microphyllus*, Sp. 38.
- 106 *M. uncinarius*, Sp. 57.
- 107 *M. trigonularis*, Sp. 56.
- 108 *M. varius*, var. Sp. 69.
- 109 *M. pinniger*, Sp. 49.
- 110 *M. ternispina*, Sp. 2.
- 111 *M. rectirostrum*, Sp. 11.
- 112 *M. formosus* Sp. 16.
- 113 *M. nigrescens*, Sp. 12.
- 114 *M. digitatus*, Sp. 77.
- 115 *M. acanthopterus?* Sp. 59.
- 116 *M. secundus*, Sp. 75.
- 117 *M. inconspicuus*, Sp. 67.
- 118 *M. Rosarium*, Sp. 87.
- 119 *M. Rota*, Sp. 73.
- 120 *M. torrefactus*, Sp. 35.
- 121 *M. torrefactus*, var. Sp. 35.

Murex

2



3



1



4



4



6



5



7

Murex

G. B. S. Junr. Exc



Murex.



16



15



17



18



19




21



20

Murex.




G.B.S. jun. exc.

Murex.

22



23




24



25



26



27



28



29

G.B.S. jun. etc.

Murex.



30



31



32



33



34



35




Murex.

36



37



38



39



40




Murex.



41




42



43



44




45

G.B.S. jun. ex.




Murex.



Murex.

54



52



53



54



Murex.

56



57



58



55

G.B.S. jun. exc.

1771



1771

Murex



59



60



61



63



62



Murex



64



65



66



67



68

Handwritten text at the top center, possibly a title or page number, which is mostly illegible due to fading.



Murex

1115



Murex


75



7L



7L



77



79



78



76

1000




Murex.

81



81



82



83




80



Small




Murex



Handwritten text, possibly a signature or title, located at the top center of the page.



Murex


Murex



94



95



91



93




96



92

1871



Murex



101



103



101



102



102


Murex



109




108



105




107



106



Murex






Murex



114



113



115



117




116



Murex

119



118



120



121

Figure 1



GENUS TYPHIS

MONTF.

A CATALOGUE

*Of all the Species hitherto known; as described by BRODERIP,
in the Zoological Proceedings, 1832;*

And figured in Part 200 of the

Conchological Illustrations,

By G. B. SOWERBY, JUN.

- 1 *T. Cumingii*, *Brod.* Bay of Caraccas, Columbia, Mr. Cuming.
- 2 *T. coronatus*, *Brod.* Salango, Columbia, Mr. Cuming.
- 3 *T. Belcheri*, *Brod.* Cape Blanco. Zool. Museum.
- 4 *T. Sowerbii*, *Brod.* Mediterranean. Philippi considers
this as identical with the fossil *T. fistulosus*,
Brocchi.
- 5 *T. pinnatus*, *Brod.*

TYPHIS.

List of Figures, Part 200.

1, 2 *T. Cumingii*.

3, 4 *T. coronatus*.

5, 6 *T. Belcheri*.

7, 8 *T. Sowerbii*.

9 *T. Sowerbii*, *var.*

10, 11 *T. pinnatus*.

Typhis

RENELIA.

Index to the Figures in parts 84, 85, 88, 89, 92, and 93, of the
CONCHOLOGICAL ILLUSTRATIONS,

BY

G. B. SOWERBY, *Jun*

Fig.

- 1, 1*. *R. pusilla*, *Brod. Z. P.* 1832, Pacific Ocean.
2. *R. pyramidalis*, *Brod. Z. P.* 1832. Panama.
3. *R. vexillum*, *Sow. Jun. Z. P.* 1841. Chiloe and Conception. The brown beaded bands are smaller and more numerous, and the outer lip more effuse than in *R. argus*.
4. *R. nitida* *Brod. Z. P.* 1832. Columbia.
- 5, 5.* *R. cruentata*, *Sow. Jun. Z. P.* 1841. Ticao, of the Philippines.
6. *R. nana*, *Sow. Jun. Z. P.* 1841. Panama.
7. *R. rugosa*, *Sow. Jun. Z. P.* 1841.
8. *R. cæolata*, *Brod. Z. P.* 1832. Panama.
9. *R. crumena*, *Lam.*
10. *R. rhodostoma*, *Beck. Sow. Jun.* 1841. Isl. Masbate, of the Philippines.
11. *R. muriciformis*, *Brod. Z. P.* 1832, Columbia.
- 11.* *R. muriciformis*, var.
12. *R. affinis*, *Brod. Z. P.* 1832. Annaa.
13. *R. tuberculata*, *Brod. Z. P.* 1832. Taheite.
14. *R. albo-fasciata*, *Sow. Jun. Z. P.* 1841. Panama, of the Philippines.

15. *R. lævigata*, Lam.
16. *R. ventricosa*, Brod. Z. P. 1832. Callao. Peru.
17. *R. elegans*, Beck. Chemn. 1270. Nicobar.
18. *R. sub-granosa*, Sow. Jun. Z. P. 1841. Bay of Manilla. This shell differs from *R. elegans* in being more finely grained and in the tubercles being much less sharp and prominent.
19. *R. pulchra*, Gray. Jay's catalogue, comonly called the "Finned Frog."
20. *R. verrucosa*, Sow. Jun. from a shell in the collection of J. Norris, Esq. resembling, in general form, *R. Rhodostona* and *R. cruentata*, but having on each side of the whorl, 3 rows, each consisting of three large, mahogany brown, smooth, rounded tubercles, between the varices, and of one on the varix. The teeth on the inner lip are interlined with red.
21. *R. scrobiculator*, var.
22. *R. neglecta*, Sow. Jun. Z. P. 1841.

Ranella.



1



2



2



3



4



4



1*

10. *P. longicauda* Lam.
 11. *P. yachay* Bond. Z. P. 1832. Callao Peru.
 12. *P. elegans* Bond. Z. P. 1832. Niadep.
 13. *P. yachay* Bond. Z. P. 1841. Bay of Man.
- The last three all differ from *P. elegans* in being more finely grained and in the tubercles being much less sharp and prominent.
14. *P. yachay*, Bond. Z. P. 1841. Bay of Man. called the "Faint Frog."
 15. *P. yachay* Bond. Z. P. 1841. Bay of Man. called the "Faint Frog."
- The last three all differ from *P. elegans* in being more finely grained and in the tubercles being much less sharp and prominent.



Ranella.



6



5



6



7



8



5



8

Handwritten text at the top center, possibly a title or page number, which is faint and difficult to decipher.



Ranella.



10



10



9



11*



11

Almond



Ranella.



15



15



12



13




14

G.B.S. jun. exc.



Ranella.


16



17




17



16



18



18






19



20



20



19



21



21



22



A CATALOGUE

OF ALL THE RECENT SPECIES HITHERTO KNOWN OF
THE GENUS

Monoceros,

OF LAMARCK,

Spec.

1. *Mon. cingulatum* *Lam.* Hist. Nat. des anim. sans vert.
Encycl. meth. t. 396. f. 4. *Sowerby*, Genera of
Recent and Fossil Shells, No. 5, *Monoceros*, f. 4.
Panama.
- 2 *Mon. imbricatum* *Lam.* l. c. Encycl. meth. t. 396. f. 1.
Sowerby, l. c. *Monoceros*, f. 1. Tierra del Fuego.
var. *Conch. Illustr.* f. 15.
- 3 *Mon. striatum* *Lam.* l. c. Encycl. meth. t. 396. f. 3. M.
breve, *Sowerby*, l. c. *Monoceros* f. 2. Tierra del
Fuego.
- 4 *Mon. glabratum* *Lam.* l. c. Encycl. meth. t. 396. f. 5.
var. *Conch. Illustr.* f. 16.
- 5 *Mon. crassilabrum* *Lam.* l. c. Encycl. meth. t. 396. f. 2.
Conch. Illustr. f. 14. var. *Conch. Illustr.* f. 13.
Valparaiso.
- 6 *Mon. cymatum*, *Sowerby*, Tank. Cat. No. 1888. Zool.
Proc. Buccm. *cymatum*, *Solander*, M. S. ined. *Monoceros*
lugubre, *Sowerby*, Genera, No. 5. *Monoceros*
f. 3. Wood Supplement. Tab. 4. *Buccinum*, f.
11-12. *Conch. Illustr.* f. 11. California.
- 7 *Mon. grande* *Gray*, in *Beechey's Narrative*. *Conch.*
Illustr. f. 1. operculum l. a. Galapagos.
- 8 *Mon. costatum*, *Sowerby*, Zool. Proc. *Conch. Illustr.* f.
6. Chili (Conception.)
- 9 *Mon. acuminatum*, *Sowerby*, Zool. Proc. *Conch. Illustr.*
f. 10. Chili (Baldivia.)

Spec.

- 10 Mon. Globulus, *Sowerby*. Zool. Proc. Conch. Illustr. f. 8. Chili (Maulé.)
- 11 Mon. citrinum, *Sowerby*, Zool. Proc. Conch. Illustr. f. 2. var. Conch. Illustr. f. 12. Coquimbo.
- 12 Mon. brevidentatum, *Gray*, in *Beechey's Narrative*. Woods Index testaceologicus, Suppl: t. 4. Buccinum. f. 10. Conch. Illustr. f. 4. Purpura cornigera, De Blainville, Annales du Museum, Monographie des Pourpres t. 9. f. 4.
- 13 Mon. punctulatum, *Gray*, in *Beechey's Narrative*. Conch. Illustr. f. 3.
- 14 Mon. unicarinum, *Sowerby*. Conch. Illustr. f. 5.
- 15 Mon. tuberculatum, *Gray*. Conch. Illustr. f. 9.
- 16 Mon. fusoides, *King*, Zool. Journ. V. p. 348. Conch. Illustr. f. 7.

CORRECTED LIST OF FIGURES

In Parts 79, 81 to 83, of the
CONCHOLOGICAL ILLUSTRATIONS.

- Fig. 1. M. grande. a operculum.
2. M. citrinum.
3. M. punctulatum.
4. M. brevi-dentatum.
5. M. unicarinum.
6. M. costatum.
7. M. fusoides.
8. M. globulus.
9. M. tuberculatum.
10. M. acuminatum.
11. M. cymatum.
12. M. citrinum. var.
13. M. crassilabrum, var.
14. M. crassilabrum.
15. M. imbricatum, var.
16. M. glabratum, var.

Monoceros*G. B. S. Jun. sc.*

Monoceros.




8



10



10



9



11





8



11



Monoceros.



1871



A CATALOGUE
OF ALL THE
RECENT SPECIES OF THE GENUS
EBURNA.

CONCHOLOGICAL ILLUSTRATIONS,

Part 20.

- 1 spirata Lam. Encycl. meth. t. 401. f. 2. E. Indies.
- 2 Ambulacrum nob. Tank. Cat. App. p. 22. Conch. Illustr.
f. 2. Java.
- 3 Valentiana Swains. Zool. Illustr. first series pl. 144. Red
Sea.
- 4 pacifica Swains. (*lutosa* Lam.) Zool. Illustr. first series pl.
146. New Zealand.
- 5 tessellata Swains. (*areolata* Lam.) Zool. Illustr. first series.
pl. 145. China, Lam.
- 6 Zeylanica Lam. Encycl. meth. t. 401. f. 3. Ceylon.
- 7 plumbea nob. Conch. Illustr. f. 3 & 4. Bucc. plumbeum
auct.^m
- 8 australis nob. (*Cancellaria spirata* Lam. secund. D. Gray.)
Conch. Illustr. f. 5. New South Wales. Shell ovate,
pinkish brown, with darker spots, spirally striated,
with a deep spiral groove close to the suture; outer
lip with a small tooth at the edge near the base.
- 9 papillaris nob. Tank. Cat. App. p. 22. Conch. Illustr. f. 1.
Cape of Good Hope.

LIST OF FIGURES.

- Fig. 1. E. papillaris.
2. E. Ambulacrum.
3 & 4. E. plumbea.
5. E. australis.

A CATALOGUE

OF THE
RECENT SPECIES OF THE GENUS
EBURNA

CONCHOLOGICAL ILLUSTRATIONS

Part 20.

1. *Eburna* Lam. Encycl. meth. t. 401. f. 2. E. Indiae.
2. *Ambulacrum* nob. Tank. Cat. App. p. 22. Conch. Illustr.
3. *E. Java*.
4. *Valentina* Swain. Noct. Illustr. first series pl. 144. Red Sea.
5. *pacifica* Swain. (Lam. Lam.) Noct. Illustr. first series pl. 145. New Zealand.
6. *remota* Swain. (Lam. Lam.) Noct. Illustr. first series pl. 146. China, Lam.
7. *Nephele* Lam. Encycl. meth. t. 401. f. 3. Cayenne.
8. *phidippus* nob. Conch. Illustr. f. 8. f. 1. Bona, phidippus nob.
9. *aurata* nob. (Conch. Illustr. f. 8. f. 1. Bona, aurata nob. D. Gray.)
10. *Conch. Illustr. f. 8. New South Wales. Shell ovate, pinkish brown, with darker spots, apically striated, with a deep spiral groove close to the suture; outer lip with a small tooth at the edge near the base.*
11. *Capensis* nob. Tank. Cat. App. p. 22. Conch. Illustr. f. 1. Cape of Good Hope.

LIST OF FIGURES

- Fig. 1. E. *pacifica*.
2. E. *Ambulacrum*.
- 3 & 4. E. *phidippus*.
5. E. *aurata*.





A CATALOGUE

OF THE

RECENT SPECIES OF CYPRÆADÆ.

In the following Catalogue the Recent Species of the family of Cypræadæ are arranged as far as the present state of knowledge will enable us according to their apparent relations. It is drawn up by Mr. G. B. SOWERBY, the Elder, who alone must be held responsible for all its imperfections, be they few or many. He has added Descriptions of such as are not already described in the Zoological Journal. As representations are given of those that are not figured before they will be referred to in this Catalogue. A few observations are added, and *no charge is made for it.*

Genus CYPRÆA.

- 1 Cypræa Princeps, *Gray*, Zool. Jour., I. p. 75, Conch. Ill. f. 1, Persian Gulf
- 2 ——— Mappa, *Linn.*, Encycl. Meth. t. 352, f. 4, *Sowerby*, Gen. of Shells, Cypræa, f. 3, Indian Ocean, Friendly Islands, Annaa, &c.
Obs. Of this beautiful species there are four or five varieties from different localities.
- 3 ——— Arabica, *Linn.*, Encycl. Meth., t. 352, f. 1, 2, Indian and Pacific Oceans.
Obs. This appears to be a variable species. The *C. Histrion*, and *C. reticulata* of some authors, Encycl. Meth., t. 351, f. 1, may possibly prove to be distinct. I have never seen the *C. maculata* of *Barnes*, but if

permitted to form my judgment from his figure (Ann. of Lyceum of Nat. Hist. of New York, vol. I., part 1, p. 133, pl. IX. f. 1.) I should say it was perfectly distinct.

4 *Cypræa Tigris*, *Linn.*, *Encycl. Meth.*, t. 353, f. 3. Indian and Pacific Oceans.

5 ——— *Pantherina*, *Solander*, *Encycl. Meth.*, t. 353, f. 5. Persian Gulf.

Obs. *C. umbilicata* of *Sowerby*, in *Tank. Cat.* may prove to be only a variety of this.

6 ——— *Mauritiana*, *Linn.*, *Encycl. Meth.* t. 350, f. 2. East Indies and Pacific.

7 ——— *stercoraria*, *Linn.*, *Encycl. Meth.* 354, f. 5. Persian Gulf.

8 ——— *Aurora*, *Solander*, *Chem. XI.*, f. 1737-1738. Ulitea, one of the Society Islands.

9 ——— *Leucodon*, *Brod.*, *Zoological Journal IV.*, t. 5. Locality unknown.

10 ——— *sulcidentata*, *Gray*, *Zool. Journ.*, t. 7 & 12, f. 5, Locality unknown.

11 ——— *Arabacula* *Lam.*, *Zool. Journ. I.*, t. 7 & 12, f. 4. St. Elena: small variety from Real Llejós.

12 ——— *Lynx*, *Linn.*, *Encycl. Meth.*, t. 355, f. 8. East Indies and Pacific.

13 ——— *Vitellus*, *Linn.*, *Encycl. Meth.*, t. 354, f. 6. East Indian Ocean, Pacific; Annaa.

14 ——— *Carneola*, *Linn.*, *Wood Ind. Test.* t. 16, f. 9. East Indian Ocean, Pacific; Annaa.

15 ——— *cinerea*, *Linn.*, *Wood Ind. Test.*, 17, f. 27. West Indies

15* ——— *Reevei*, *Gray*. Shell oval, ventricose, brown; spire evident, aperture narrow, ends short, reddish; teeth

small, numerous, continuing over the columella and strongly marked over the cavity at its anterior extremity; length 1.2, breadth 0.85. From Garden Island, mouth of the Swan River. Conch. Illustr., f. 52.

- 16 *Cypræa melanostoma*, *Leathes*, Zool. Journ., t. 18, f. 3, 4. Persian Gulf.

- 17 ——— *achatina*, *Solander*, Wood Ind. Test., t. 17, f. 18. Annaa, South Seas

- 18 ——— *arenosa*, *Gray*, Zool. Journ., I. t. 7 & 12, f. 6. Pacific Ocean, Annaa.

- 19 ——— *nivosa*, *Brod.*, Zool. Journ. III., t. 4, f. 1, Mauritius

- 20 ——— *Broderipii*, *Gray*, Conch. Illus., f. 2. Madagascar.

Shell oval, ventricose, rather acuminate in front, of a pale fulvous colour varied with rosy hue and whitish, and for the most part covered with variously sized brown reticulations, dorsal line rather lateral, wide, not covered with the brown reticulations; aperture rather narrow, teeth nearly covering the whole of the columellar lip, white, rather large, those of the columellar lip smaller; base orange brown; sides darker, cracked, with some white spots about their upper edge; extremities brown: length 2.65, breadth 1.75, Conch. Illust. f. 2.

C. Broderipii, *Gray*, Descrip. Cat. p. 3, no. 18.

Obs. The only specimen known is in Mr. Cummings Cabinet.

- 21 ——— *Exanthema*, *Linn.*, Encycl. Meth., t. 349. West Indies.

- 22 ——— *Cervus*, *Linn.*, Encycl. Meth., t. 351, f. 3. East Indies

- 23 *Cypræa Testudinaria*, *Linn.*, *Encycl. Meth.*, t. 351, f. 2, Mauritius.
- 24 ——— *Talpa*, *Linn.*, *Encycl. Meth.*, t. 353, f. 4, Indian Ocean, Annaa.
- 25 ——— *exusta*, *Conch. Illustr.*, f. 2*. Persian Gulf.
 Shell oblong-oval, dark brown, beneath pitch black; extremities rather produced; teeth very numerous, close-set, small, short, particularly those of the columellar lip; base of the columella white, with two rather obsolete teeth near the anterior extremity; teeth dark brown, not extending over the columella, nor over the concavity near its base; space between the teeth and the fold forming the anterior canal very narrow. Length 2.6, breadth 1.4. Inhabits the Persian Gulph.
Obs. This is the *Cypræa Talpa exustus* of Gray's *Descr. Catal.* p. 4, no. 28. It appears to be quite distinct.
- 26 ——— *Argus*, *Linn.*, *Encycl. Method.*, 350, f. 1. Indian Ocean; Friendly Islands.
- 27 ——— *Scurra*, *Chem.*, *Encycl. Meth.*, t. 352, f. 3. Annaa, East Indies.
- 28 ——— *pulchra*, *Gray*, *Zool. Journ.*, Vol. I. t. 7. and 12, f. 9. Persian Gulf.
- 29 ——— *Isabella*, *Linn.* *Wood's Cat.*, t. 17, f. 18. East Indian and Pacific; Annaa.
- 30 ——— *controversa*, *Gray*, *Zool. Journ.*, t. 7 & 12, f. 7.
Obs. This may prove to be only a variety of *C. Isabella*
- 31 ——— *lurida*, *Linn.*, *Encycl. Meth.*, t. 354, f. 2. Mediterranean.
- 32 ——— *microdon*, *Gray*, *Conch. Illust.* f. 3.

- 33 *Cypræa Scottii* *Brod.*, Zool. Jour., Vol V., p. 330, t. XIV. Garden Island.
- 34 ——— Mus, *Linn.*, Encycl. Meth., t. 354, f. 1.
Obs. The *Ovula tuberculosa*, *Duclos*, is a decided *Cypræa*, and should come next to this; it is the same as Mr. Gray has called *Cypræa Deshayesii*.
- 35 ——— tessellata, *Swainson*, Zool. Illust., t. 111. South Sea Islands.
- 36 ——— annulata, *Gray*, Conch. Illust. f. 4. Annaa.
- 37 ——— Margarita, *Sol.*, Conch. Illust., f. 6. Ulitea.
- 38 ——— Cicercula, *Linn.*, Encycl. Meth., t. 355, f. 1. Annaa.
- 39 ——— Globulus, *Linn.*, Encycl. Meth., t. 356, f. 2, East Indies
- 40 ——— Staphylæa, *Linn.*, Encycl. Meth., t. 356, f. 9. Mauritius.
- 41 ——— Nucleus, *Linn.*, Encycl. Meth. t. 355, f. 3. Mauritius; Annaa.
- 42' ——— Madagascariensis, *Lam.*, Sowerby's Genera, *Cypræa*, f. 6. Madagascar, Pacific.
- 43 ——— guttata, *Gmel.*, Zool. Journ., II., t. 18, f. 1, 2.
- 44 ——— poraria, *Lam.*, Encycl. Meth., t. 356, f. 4, Ceylon; Pacific.
- 45 ——— albuginosa, *Gray*, Zool. Journ., I., t. 7 and 12, f. 2, Ceylon; Mexico.
- 46 ——— Listeri, *Gray*, Lister, t. 701, f. 48.
- 47 ——— gangrenosa, *Gmel.*, Conch. Illustr., f. 8, Ceylon
- 48 ——— citrina, *Gray*, Conch. Illustr., f. 9.
- 49 ——— Helvola, *Linn.*, Encycl. Meth., t. 356, f. 13. Tahiti.

- 50 *Cypræa bicallosa*, *Gray*, *Conch. Illustr.*, f. 10.
Shell ovate-oblong, extremities rather produced, margined, orange brown, back paler with brown dots; a light coloured callosity at each end of the back; base darker coloured; teeth large, close, much extended; length breadth
- 51 ——— *spurca*, *Linn.*, *Encycl. Meth.*, t. 356, f. 14.
var. *alba*, *Conch. Illustr.*, f. 53.
- 52 ——— *flaveola*, *Linn.*, *Conch. Illustr.*, f. 11.
- 53 ——— *erosa*, *Linn.*, *Encycl. Meth.* t. 355, f. 4.
Annaa.
- 54 ——— *Lamarckii*, *Gray*, *Conch. Illustr.*, f. 12.
- 55 ——— *ocellata*, *Linn.*, *Encycl. Meth.*, t. 355, f. 7.
Ceylon.
- 56 ——— *Turdus*, *Lam.*, *Zool. Journ. I.*, t. 7 and 12, f. 3.
var. *alba*, *Conch. Illustr.*, f. 54.
- 57 ——— *Caput-serpentis*, *Linn.*, *Encycl. Meth.*, t. 354, f. 4. Tahiti,
- 58 ——— *Annulus*, *Linn.*, *Encycl. Meth.*, t. 356, f. 7.
Tahiti.
- 59 ——— *obvelata*, *Lam.*, *Conch. Illustr.*, f. 13. Tahiti.
- 60 ——— *Moneta*, *Linn.*, *Encycl. Meth.*, t. 356, f. 3.
- 61 ——— *Asellus*, *Linn.*, *Encycl. Meth.*, t. 356, f. 5.
- 62 ——— *4-maculata*, *Gray*, *Wood's Cat. App.*, t. 3. f. 8.
- 63 ——— *interrupta*, *Gray*, *Conch. Illustr.*, f. 15, Ceylon.
- 64 ——— *Hirundo*, *Linn.*, *Wood's Cat.*, t. 17., f. 36,
var. *Owenii*, *Conch. Illustr.*, f. 12**.
- 65 ——— *stolida*, *Linn.*, *Chemn.*, XI., 1743, 1744,
Ceylon.
- 66 ——— *neglecta nobis*, *Conch. Illustr.*, f. 12*.

Shell ovate-oblong, bluish; with an irregular rust-colored blotch in the middle of the back, ends produced,

spire depressed ; two black dots at each end ; margins and base white, the former brown dotted : teeth numerous, much extended on both sides, length 0·7., breadth 0·4. From the Mauritius.

C. Hirundo, var. 1. *Gray*, Descr. Cat., p. 9. *C. stolid*a, var. nobis.

Obs. A very pretty little shell, which may be distinguished at first sight with ease from both *C. Hirundo* and *C. stolid*a, though more like the latter than the former.

67 *Cypræa punctata*, *Linn.*, Encycl. Meth., t. 355, f. 10, Mauritius.

68 ——— *pulchella*, *Swainson*, Conch. Illustr., f. 40.

69 ——— *pyriformis*, *Gray*, Conch. Illustr., f. 23, Ceylon.

70 ——— *Walker*i, *Gray*, Conch. Illustr., f. 22*.

Shell ovate, somewhat pear-shaped, pale with yellow brown dots, and a broad interrupted central band ; spire depressed ; base and margin yellowish, the latter with a few purple dots ; anterior extremity with two purple spots, posterior with an irregular dark blotch ; teeth whitish, their interstices purple, those of the inner lip small, close-set, and somewhat extended ; inside purple ; length 0·9. breadth 0·5.

*Cypræa Walker*i, *Gray*, Desc. Cat., *Cypræadæ*, p. 11.

Hab. in the Persian Gulph. Collection of Mr. Walker.

71 ——— *Humphreys*i, *Gray*, Zool. Journ., I. p. 489, Conch. Illust., f. 55.

72 ——— *teres*, *Gmel.*, Conch. Illust., f. 56.

Shell oblong, cylindrical, pale purplish brown ; with three interrupted bands, consisting of short longitudinal stripes, the middle band broadest ; spire umbilicated ; base white ; margin white with a few brown spots ;

extremities produced, turned upwards; outer lip rather thickened, reflected, with its front depressed; teeth small, numerous, close-set, all within the aperture, which is narrow and linear, length 0·8. breadth 0·35.

From Haynam. This species approaches nearest to *C. tabescens* of which Mr. Gray considers it a variety: it appears to me to be easily distinguished by the above characters.

73 ——— *tabescens*, *Solander*, *Conch. Illustr.*, f. 14, Ceylon, Tahiti.

74 ——— *cylindrica*, *Born.*, *Sowerby's Genera*, *Cypræa*, f. 4. Tahiti.

75 ——— *Cribraria*, *Linn.*, *Encycl. Meth.*, t. 355, f. 5, Mauritius, var. *Conch. Illustr.*, f. 16*.

76 ——— *Goodalli*, *Gray*, *Conch. Illustr.*, f. 16.

Shell oblong, white, margins with a few brown dots; back with a nearly central irregular brown blotch, having a few white spots on it; spire depressed; teeth small; length 0·55, breadth 0·3. Found by Mr. Cuming. Lord Hood's Island.* *C. Goodallii* Gray *Descrip. Cat. Cypræadæ*, p. 10.

77 *Cypræa Cumingii*, *Gray*, *Conch. Illustr.*, f. 5.

Shell oblong, ends produced, spire depressed; back pale fulvous, with round white spots, margins with round very dark brown dots, outer margin thickened; base white, mouth long and narrow, teeth very small: length 0·6. breadth 0·3. *Cypræa Cumingii*, Gray, *Descr. Cat.*, p. 6, No. 41.

Obs. Two specimens were found by Mr. Cuming.

78 ——— *Caurica*, *Linn.*, *Encycl. Meth.*, t. 356, f. 10.

* *Lord Hood's Island* and *Annaa* or *Chain Island* belong to the Paumotian group.

- 79 ——— cruentata, *Gmel.*, *Encycl. Meth.*, t. 353, f. 2.
 80 ——— erronea, *Linn.*, *Wood's Cat.* t. 17, f. 39.
 81 ——— felina, *Gmel.*, *Wood's Cat.* t. 17, f. 26.
 82 ——— irrorata, *Solander*, *Conch. Illust.* f. 25. Elizabeth Island.
 83 ——— Spadicea, *Swains.*, *Zool. Illust.* t. 182. Upper and lower figures.
 84 ——— Onyx, *Linn.*, *Wood's Cat.* t. 17, f. 31.

Obs. There are several varieties of this species, one of which is *C. adusta* of Lam. : and another very beautiful variety of a fine flesh-color is represented at f. 17 of these *Conch. Illust.*

- 85 ——— Pyrum, *Gmel.*, *Encycl. Meth.* t. 373, f. 1. Mediterranean.
 86 ——— picta, *Gray*, *Zool. Journ.* t. 7 & 12, f. 10. Guaymas.
 87 ——— zonata, *Lam.*, *Zool. Journ.*, t. 7 & 12, f. 8.
 88 ——— Xanthodon, *Gray*, *Conch. Illust.* f. 18.

Shell ovate, bluish brown, with central and posterior darker transverse bands and dotted with olive brown; base rounded, fulvous, sides rather thickened, lighter, with a few large black spots; teeth reddish brown; inside dark purple: length 1.2, breadth 0.75. *Cypræa* Xanthodon, *Gray's Descrip. Cat.*, p. 10.

- 89 ——— nigro-punctata, *Gray*, *Conch. Illust.* f. 22. Gallapagos Islands.
 90 ——— pallida, *Gray*, *Conch. Illust.*, f. 19.
 91 ——— punctulata, *Gray*, *Conch. Illust.*, f. 23. Panama.
 92 ——— ziczac, *Linn.*, *Encycl. Meth.*, t. 356, f. 8.
 93 ——— undata, *Solander*, *Wood's Cat.*, t. 17, f. 23.
 94 ——— clandestina, *Linn.*, *Wood's Cat.*, app. t. 3, f. 17.

- 95 *Cypræa lentiginosa*, *Gray*, Zool. Journ., t. 7 & 12, f. 1. Ceylon.
- 96 ——— *contaminata*, *Gray*, Conch. Illust., f. 21.
 Shell pale buff colour, oblong pear-shaped; base and margin with dark brown dots; dorsal margin of the outer lip slightly reflected; spire depressed, extremities produced; teeth small, close set. Length 0.5, breadth 0.27. *C. contaminata*, *Gray's Descriptive Cat.* p. 11, No. 88.
- 97 ——— *sanguinolenta*, *Gmel.*, Swains. Zool. Illust. t. 182, middle figures.
- 98 ——— *fimbriata*, *Dillw.* Wood's Cat., t. 17, f. 57. Sibo, Mauritius, &c.
- 99 ——— *angustata*, *Gmel.*, Wood's Cat., t. 17, f. 32. South Africa.
- 100 ——— *piperita*, *Gray*, Conch. Illust., f. 24. New South Wales.
- 101 ——— *Algoensis*, *Gray*, Conch. Illust., f. 26. Southern Africa.
- 102 ——— *edentula, nobis*, Conch. Illust., f. 26.* Southern Africa

Shell pear-shaped, fulvous, dotted with brown; beneath and border lighter, with darker and larger dots; margin of the outer lip thickened; aperture rather wide, without teeth; anterior extremity of the columella curved downwards, without any folds; spire distinct, flat; posterior extremity rather produced; length 0.9, breadth 0.6.

Obs. This, according to Mr. Gray, is a variety of his *Luponia Algoensis*, "with the teeth more or less obliterated:" but there never have been the slightest vestige of teeth in any of the specimens we have seen,

nor are there any irregular ridges crossing the front edge of the columella, which form the distinguishing mark of his genus *Luponia*. It is an extremely rare and very distinct species from Southern Africa.

- 103 *Cypræa similis*, *Gray*, *Conch. Illust.*, f. 27. Cape of Good Hope.

Shell pear-shaped, smooth, pale brown, with the border and base lighter and spotless: anterior end of the columella crossed by two or three irregular ridges; teeth of the outer lip large, distant; those of the inner lip closer, slightly extended: length 1.4, breadth 0.9.

C. similis, *Gray*, *Zool. Misc.* p. 36.

- 104 ——— *fusco-dentata*, *Gray*, *Conch. Illust.*, f. 28. Cape of Good Hope

- 105 ——— *capensis*, *Gray*, *Wood's Cat.*, app. t. 3, f. 18. Cape of Good Hope.

- 106 ——— *pustulata*, *Lam.*, *Sowerby's Genera*, f. 5. Panama and Isle of Plata, Mexico.

- 107 ——— *Adamsonii*, *Gray*, *Conch. Illust.*, f. 7. Mauritius.

Shell pearshaped, with numerous transverse ribs, decussated by less distinct and more distant longitudinal ribs; whitish, mottled with pale brown; teeth numerous, close-set and sharp: length 0.4, breadth 0.25.

C. Adamsonii, *Gray*, *Descrip. Cat. of Shells*, p. 7, *Cypræadæ* No. 48.

- 108 ——— *ovulata*, *Lam.*, *Encycl. Meth.*, t. 355, f. 2, Cape of Good Hope

- 109 ——— *aperta*, *Swainson*, *Wood's Cat.*, app. t. 3, f. 10. Cape of Good Hope.

- 110 ——— *carnea*, *Gray*, *Wood's Cat.*, app. t. 3, f. 15.

- 111 *Cypræa Maugeriæ*, *Gray*, *Conch. Illust.*, f. 30. Gallapagos.
 Shell oval, thin, pellucid, rose coloured, with darker ends and a central dorsal blotch; ribs thin, numerous, rather close, the interstices roughish; dorsal groove distinct, narrow; lips whitish, teeth sharp: length 0·5, breadth 0·35. *Cypræa Maugeriæ*, *Gray*, *Descr. Cat.* p. 13, No. 111.
- 112 ——— *Australis*, *Lam.*, *Conch. Illust.*, f. 29. New South Wales.
- 113 ——— *rufescens*, *Gray*, *Conch. Illust.*, f. 31.
 Shell nearly globular, pale reddish, thin, pellucid, ribs thin, rather close; under side white, outer lip rather narrow; no dorsal line; length 0·4, breadth 0·3. *Cypræa rufescens*, *Gray*, *Descr. Cat.* p. 13, No. 112.
- 114 ——— *Europæa*, *Lam.*, *Encycl. Meth.*, t. 356, f. 1. b.
- 115 ——— *sanguinea*, *Gray*, *Conch. Illust.*, f. 32. Panama and Mexico.
 Shell ovate, dark purplish brown, with a dark crimson blotch on the back, dorsal line short, its ends whitish; ribs rather distant, smooth, lighter coloured: length 0·45, breadth 0·30. *Cypræa sanguinea*, *Gray*, *Descr. Cat.* p. 14, No. 119.
- 116 ——— *quadripunctata*, *Gray*, *Conch. Illust.*, f. 33.
- 117 ——— *globosa*, *Gray*, *Conch. Illust.*, f. 34.
 Shell globular, reddish, closely and acutely ribbed; dorsal line narrow, distinct, running from the posterior to the anterior extremity; margin thickened; under side rounded, white; teeth small, sharp: length 0·3, breadth 0·25. *Cypræa globosa*, *Gray*, *Descr. Cat.* p. 14, No. 121.
- 118 ——— *exigua*, *Gray*, *Conch. Illust.*, f. 35.
 Shell ovate, pellucid, white, varied with rose colour;

extremities subrostrated; dorsal ribs thick, distant; lateral and ventral ribs thin and close, and teeth small; no dorsal groove: length 0·2, breadth 0·15. New South Wales, G. Humphrey. *Cypræa exigua*, Gray, Descr. Cat. p. 15, No. 123.

119 *Cypræa subrostrata*, Gray, Conch. Illust. f. 36.

120 ——— *fusca*, Gray, Conch. Illust., f. 37. Gallapagos and Bay of Guyaquil. Mus. Cuming.

Shell ovato-globose, dark brown, with a pale dorsal streak, extremities slightly produced; ribs small, rather close, some of them not continued to the edge of the lips beneath: length 0·33, breadth 0·25. *Cypræa fusca*, Gray, Descr. Cat. p. 15, No. 125.

121 ——— *Oryza*, Lam., Conch. Illust., f. 38.,

122 ——— *nivea*, Gray, Conch. Illust., f. 38*.

123 ——— *Pulex*, Solander, Conch. Illust., f. 32,

124 ——— *Pediculus*, Linn, Encycl. Meth., t. 356, f. 1, a.

125 ——— *Pacifica*, Gray, Conch. Illust., f. 39*. Gallapagos.

Shell ovato-oblong, brown, with minute dots and four squarish darker brown alternating spots on the sides of the narrow dorsal groove; ribs narrow, rather close set; sides rather thickened, rose coloured, base and teeth white: length 0·4, breadth 0·25.

Cypræa pacifica, Gray, Descr. Cat. p. 15, No. 133.

126 ——— *suffusa*, Gray, Conch. Illust., f. 41.

Shell ovate, rose-coloured, dotted and varied with reddish brown, base paler; dorsal groove narrow, ribs close-set, granose on the back; anterior end on each side of the canal rose-red: length 0·45, breadth 0·30.

Cypræa suffusa, Gray, Descr. Cat. p. 16, No. 134.

127 ——— *Californica*, Gray, Conch. Illust., f. 42.

- 128 *Cypræa Solandri*, *Gray*, *Conch. Illust.*, f. 43.

Shell reddish brown, with a light-coloured dorsal groove, and a dark streak along each side of it; ribs prominent, wide apart, tuberculated at their dorsal extremities, mouth rather narrow, teeth of the outer lip more numerous than in *C. radians*: margin generally broad: length 0·7, breadth 0·55. Pacific Ocean.

Obs. This species is very much like *radians*, indeed it is impossible to know which of the two, this or the one we commonly call by that name, is the *radians* of Lam. The principal difference is in the number of teeth on the inner edge of the outer lip.

- 129 ——— *radians*, *Lam.*, *Wood's Cat.* t. 17, f. 58.

- 130 ——— *depauperata*, *Nobis*, *Conch. Illust.*, f. 49.

Shell nearly orbicular, rather depressed. pale fawn coloured, dorsal line narrow, distinct; transverse ribs few, distant, sometimes interrupted; margins thickened; under side somewhat rounded; ribs and teeth of the outer lip more numerous and closer than those of the inner: length 0·5, breadth 0·4.

The locality of this species, which does not appear to be nearly related to any other is unknown; very few specimens have occurred.

Genus, *ERATO*, *Gray*.

Obs. I readily adopt this Genus, which appears to have been first established by Risso, followed by Gray: merely observing that it is a link connecting *Cypræa* with *Marginella*.

- 1 *Erato scabriuscula*, *Gray* : *Marginella Cypræola*, *Sowerby*, *Conch. Illust.* f. 45.

Shell ovate, turbinate, purplish brown, covered, except the dorsal line, with minute granules ; spire short, conical ; aperture rather wide, paler coloured ; columellar lip plaited transversely its whole length ; outer lip thickened, largely toothed on its inner edge ; anterior canal wide : length 0·4, breadth 0·2. Brought by Mr. Cuming from St. Elena and Acapulco.

Obs. Young shell destitute of the minutely granular coat, and of course not showing the dorsal line.

- 2 — *sulcifera*, *Gray*, *Conch. Illustr.* f. 46.

Shell ovate, turbinate, pallid, very minutely and indistinctly granular ; dorsal line wide, deeply impressed ; aperture linear, very narrow, outer lip thickened, nearly white, minutely toothed on its inner margin ; anterior part of the columellar lip minutely toothed ; a single red spot at the anterior extremity : length 0·2, breadth 0·1. From the Cape of Good Hope ; a single specimen in the late G. Humphrey's collection.

- 3 — *lævis*, *Gray* ; *Cypræa Voluta* *Mont. Test. Brit.* p. 203, t. 6, f. 7. *Voluta lævis*, *Dowson*. Coasts of Britain and Mediterranean, *Conch. Illustr.* f. 57.

- 4 — *Maugeriæ*, *Gray*, *Conch. Illustr.* f. 47.

Shell somewhat conical, turbinate, livid ; spire very short, last volution ventricose, very obtusely angular behind ; outer lip thickened, nearly white, toothed along its inner margin ; a few very indistinct teeth near the front of the columella : length 0·2, breadth 0·15. West Indies, common.

- 5 — *Lachryma*, *Gray*. *Lachryma trifasciata*, *Humphrey*, *MSS.*, *Conch. Illust.* f. 48.

Shell oval, turbinate, whitish, with three red transverse

bands, smooth; aperture narrow; outer lip somewhat thickened, minutely toothed on its inner edge; columellar lip slightly concave, toothed anteriorly: length 0.2, breadth 0.13. From New South Wales; several specimens were in the cabinet of the late G. Humphrey.

6 *Erato Guttula*, *Sowerby*, *Conch. Illustr.* f. 50.

Shell ovate, rather ventricose, cream coloured; spire very short, indistinct, aperture narrow, rather effuse anteriorly; outer lip thickened, very minutely and indistinctly toothed along its inner margin; columellar lip with three folds anteriorly: length 0.25, breadth 0.15. From the Mauritius.

Obs. When worn the spire is distinct.

7 ——— *angistoma*, *Sowerby*, *Semi-Cowry*, *G. Humphrey*.

Shell ovate, gibbose, pallid; spire short; aperture as long as the whole shell, very narrow; outer lip rather thickened, wide, very minutely toothed along its inner margin; columella with six or seven anterior folds and internal concavity: length 0.2, breadth 0.15.

From the East Indies. *Conch. Ill.* f. 51.

Genus OVULUM.

- | | |
|---|---|
| 1 | <i>Ovulum Ovum</i> , <i>Sowerby</i> , <i>Spec. Conch. Ovulum</i> , f. 1 to 5. |
| 2 | ———— <i>Margarita</i> „ „ f. 19, 20. |
| 3 | ———— <i>Adriaticum</i> „ „ f. 23, 24. |
| 4 | ———— <i>Pyriforme</i> „ „ f. 21, 22, 25. |
| 5 | ———— <i>Carneum</i> „ „ f. 17, 18. |
| 6 | ———— <i>Marginatum</i> „ „ f. 15, 16. |
| 7 | ———— <i>Lacteum</i> „ „ f. 13, 14. |
| 8 | ———— <i>Breve</i> „ „ f. 26, 27. |
| 9 | ———— <i>verrucosum</i> „ „ f. 10, 11, 12. |

- | | |
|----|--|
| 10 | Ovulum angulosum, <i>Sowerby, Sp. Con. Ovulum</i> f. 6 to 9. |
| 11 | ——— triticeum " " f. 35. |
| 12 | ——— Frumentum " " f. 37. |
| 13 | ——— striatulum " " f. 38. |
| 14 | ——— emarginatum " " f. 54, 55. |
| 15 | ——— gibbosum " " f. 28 to 31. |
| 16 | ——— obtusum " " f. 34. |
| 17 | ——— Seminulum " " f. 40. |
| 18 | ——— Formicarium " " f. 39. |
| 19 | ——— Secale " " f. 36. |
| 20 | ——— Spelta " " f. 43. |
| 21 | ——— intermedium " " f. 32, 33. |
| 22 | ——— birostre " " f. 41, 42, 44, 45. |
| 23 | ——— longirostratum " " f. 46 to 48. |
| 24 | ——— Volva " " f. 56, 57. |
| 25 | ——— aciculare " " f. 49 to 52. |
| 26 | ——— patulum " " f. 58. |
| 27 | ——— hordeaceum " " f. 53. |
| 28 | ——— rufum. Shell oblong, acuminated at the posterior extremity, red; outer lip thickened, paler; aperture narrow, broader anteriorly; columella with a depressed longitudinal line within, and a slightly spiral posterior plait; length 0.5, breadth 0.15. Varies in being wholly of a pale colour. A few specimens were dredged in sandy mud at a depth of seven fathoms in the Bay of Caraccas by Mr. Cuming. <i>Conch. Ill. Cyprædæ</i> , f. 58. |
| 29 | ——— Avena. Shell oblong, red, extremities rather acuminated, back somewhat gibbous, very finely transversely striated: outer lip thickened, paler; aperture narrow, broader at the anterior, and notched at the posterior extremity; columella with a single plait poste- |

riorly; length 0.55. breadth 0.22. Found by Mr. Cuming at Conchagua in Mexico. Varies in the intensity of its colouring. *Conch. Ill. Cypræadæ*, f. 59.

30 *Ovulum inflexum*. Shell oblong, somewhat cylindrical, smooth, pale; posterior extremity somewhat beaked, beak inflected; aperture rather effuse anteriorly; outer lip thickened, columellar lip keeled within, with a single plait posteriorly, length 0.7. breadth 0.25. A single specimen was found by Mr. Cuming in the Gulph of Dulce. Two were in the late G. Humphrey's Collection. *Conch. Ill. Cypræadæ*, f. 60.

31 ——— *æquale*. Shell oblong, somewhat cylindrical, red; outer lip thickened; extremities rather obtuse; aperture rather broad, equal at each extremity; columella with a distinct internal keel, length 0.45. breadth 0.18. Found at Panama by Mr. Cuming. *Conch. Ill. Cypræadæ*, f. 61.

INDEX

TO

THE CYPRÆIDÆ,

Contained in parts 1 to 8, and 101 to 131, inclusive.

*** The numbers after the names indicate the situation of the species in the Catalogue.

FIG

- 1 *Cypræa Princeps*, 1
- 2 — *Broaderipii*, 22
- 2* — *exusta*, 35
- 3 — *microdon*, 32
- 4 — *annulata*, 36
- 5 — *Cumingii*, 77
- 6 — *Margarita*, 37
- 7 — *Adamsonii*, 107
- 8 — *gangrenosa*, 47
- 8* — *gangrenosa*, var.
- 9 — *citrina*, 48
- 10 — *bicallosa*, 50
- 11 — *flaveola*, 52
- 12 — *Lamarckii* and var., 54
- 12* — *Stolida*, var., 65
- 12** — *Hirundo*, var., 64
- 13 — *obvelata*, 59
- 14 — *tabescens*, 73
- 15 — *interrupta*, 63
- 16 — *Goodallii*, 76
- 16* — *cribraria*, var., 75
- 17 — *Onyx*, var., 84
- 18 — *Xanthodon*, 88
- 19 — *pallida*, 90
- 20 — *punctulata*, 20
- 21 — *contaminata*, 96
- 22 — *nigro-punctata*, 89
- 22* — *Walkeri*, 70
- 23 — *pyriformis*, 69
- 24 — *piperita*, 100
- 25 — *irrorara*, 82
- 26 — *Algoensis*, 101
- 26* — *edentula*, 102
- 27 — *similis*, 103
- 28 — *fusco-dentata*, 104
- 29 — *australis*, 112
- 30 — *Maugeriæ*, 111
- 31 — *rufescens*, 113
- 32 — *sanguinea*, 115
- 32* — *Pulex*, 123
- 33 — *4-punctata*, 116
- 34 — *globosa*, 117
- 35 — *exigua*, 118
- 36 — *subrostrata*, 119

F G.

- 37 *Cypræa fusca*, 120
- 38 — *Oryza*, 121
- 38* — *nivea*, 122
- 39* — *pacifica*, 125
- 40 — *pulchella*, 28
- 41 — *suffusa*, 126
- 42 — *Californica*, 127
- 43 — *Solandri*, 128
- 44 — *Scottii*, 33
- 45 *Erato scabriuscula*
- 46 — *sulcifera*
- 47 — *Maugeriæ*
- 48 — *Lachryma*
- 49 *Cypræa depauperata*, 130
- 50 *Erato Guttula*
- 51 — *angistoma*
- 52 *Cypræa Reevii*, 15*
- 53 — *Spurca*, var. *alba*, 51
- 54 — *Turdus*, var. *alba*, 56
- 55 — *Humphreysii*, 71
- 56 — *teres*, 72
- 57 *Erato lævis*
- 58 *Ovulum rufum*
- 59 — *Avena*
- 60 — *inflexum*
- 61 — *æquale*
- 63 *Cypræa cribraria*, 75
- 64 — *melanostoma*, 16
- 65 — *melanostoma*, var.
- 66 — *Vitellus*, 13
- 67 — *ocellata*, 55
- 68 — *poraria*, 44
- 69 — *Childreni*
- 70 — *Mappa*, var. S. Pacific
- 71 — *pustulata*, 106 [Ocean. 2]
- 72 — *Pyrum*, 85
- 73 — *Achatina*, 17
- 74 — *pulchella*, 68
- 75 — *arenosa*, 18
- 76 — *pallida*, 90
- 77 — *arabacula*, 11
- 78 — *Globulus*, 39
- 79 — *zonota*, 87
- 80 — *Histro.* (*Arabica* 3)

INDEX.

FIG		FIG	
81	<i>Cypræa spurca</i> , 51	135	<i>Cypræa Felina</i> , var., 81
82	— <i>lurida</i> , 31	136	— <i>controversa</i> , 30
83	— <i>staphylæa</i> , 40	137	— <i>Felina</i> , 81
84	— <i>Cicercula</i> , 38	138	— <i>fimbriata</i> , var., 98,
85	— <i>arabica</i> , 3	139	— <i>lentiginosa</i> , 95
86	— <i>Nucleus</i> , 41	140	— <i>fimbriata</i> , 98
87	— <i>clandestina</i> , 94	141	— <i>Aurora</i> , 8
88	— <i>4-maculata</i> , 62	142	— <i>europæa</i> , 114
89	— var	142*	— <i>europæa</i> , var., 114
90	— <i>Tigris</i> , 4	143	— <i>Ziczac</i> , 92
91	— <i>stolida</i> , 65	144	— <i>aperta</i> , 109
92	— var	145	— <i>ovulata</i> , 108
93	— <i>Asellus</i> , 61	146	— <i>radians</i> , 129
94	— <i>tessellata</i> , 55	147	— <i>carnea</i> , 110
95	— <i>spadicea</i> , 83	148	— <i>Pediculus</i> , 124
96	— <i>Lamarckii</i> , var. <i>alba</i> ., 54	149	— <i>Candidula</i> . <i>Gask.</i> <i>Zool.</i>
97	— <i>Beckii</i> , <i>Gask.</i>		<i>Proc.</i> 1835
98	— <i>Isabella</i> , 29	150	— <i>rubinicolor</i> . <i>Gask.</i> ditto
99	— <i>Mappa</i> , var., 2	151	— <i>formosa</i> . <i>Gask.</i> ditto
100	— <i>nivosa</i> , 19	152	— <i>testudinaria</i> , 23
101	— <i>cylindrica</i> , 74	153	— <i>labiosa</i> . <i>Gask.</i> <i>Zool.</i> <i>Proc.</i>
102	— <i>albuginosa</i> , 45		1836
103	— <i>Indica</i> , (<i>Scurra</i> , 27)	154	— <i>vesicularis</i> . <i>Gask.</i> ditto
104	— <i>spurca</i> , 51	155	— <i>producta</i> . <i>Gask.</i> ditto
105	— <i>angustata</i> , 99	156	— <i>Mus.</i> , 34
106	— <i>Scurra</i> , 27	157	— <i>Mus.</i> var?
107	— <i>Lynx</i> , 12	158	— <i>Caurica</i> , 78
108	— <i>sanguinolenta</i> , 97	159	— <i>Caurica</i> , var.
109	— <i>undata</i> , 93	160	— <i>Caurica</i> , var.
110	— <i>sulcidentata</i> , 10	161	— <i>erythræensis</i> . <i>Beck</i>
111	— <i>Capensis</i> , 105	162	— <i>picta</i> , 86
112	— <i>cruentata</i> , 79	163	— <i>cinerea</i> , 15
113	— <i>Talpa</i> , 24	164	— <i>Mauritiana</i> , 6
114	— <i>Listeri</i> , 46	165	— <i>carneola</i> , 14
115	— <i>Annulus</i> , 58	166	— <i>arabica</i> . <i>Var. riticulata</i> . 3
116	— <i>Madagascariensis</i> , 42	167	— <i>stercoraria</i> , 7
117	— <i>punctulata</i> , 91	168	— <i>Pantherina</i> , 5
118	— <i>Lynx</i> , var., 12	169	— <i>Pantherina</i> . <i>Var umbili-</i>
119	— <i>erosa</i> , var., 53		<i>cala</i> , 5
120	— <i>vesicularis</i> . <i>Gask</i>	170	— <i>exanthema</i> , 21
121	— <i>Helvola</i> , 49	171	— <i>erosa</i> , 53
122	— <i>citrina</i> , 48	172	— <i>erosa</i> , var.
123	— <i>Moneta</i> , 60	173	— <i>Turdus</i> , 56
124	— <i>errones</i> , var., 80	174	— <i>Hirundo</i> , 64
125	— <i>Argus</i> , 26	175	— <i>Cervus</i> , 22
126	— <i>pulchra</i> , 28	176	— <i>guttata</i> , 43
127	— <i>Caput-Serpentis</i> , var., 57	177	— <i>Leucodon</i> , 9. Side view
128	— <i>errones</i> , var., 80		Part 131
129	— <i>errones</i> , 80	178	— <i>Leucodon</i> . Front view.
130	— <i>Moneta</i> , var., 60	179	— <i>achatidea</i> . <i>Gray</i>
131	— <i>Caput-Serpentis</i> , 57	180	— <i>errones</i> , var., 80 (<i>C. sub-</i>
132	— <i>errones</i> , var., 80		<i>viridis</i> . <i>Reeve</i>)
133	— <i>adusta</i> (<i>Onyx</i> 84)	187	— <i>Cumingii</i> , var., 77
134	— <i>Pantherina</i> , var., 5		

N.B. Parts 1 to 8, and 101 to 131, inclusive, will be found to contain a complete illustrated monograph of the Genus *Cypræa*.


The List published with each separate Part may now be cancelled.

15th, Nov, 1837.

Cypræa.



G. B. S. exc.



Cypræa.





Cypræa.





Cypræa.



22



25



21



21



2



12 xx



12 xx



24




23




23

Plates



Cypræa.





Cypræa.

32^m






Cypræa.




39*



Cypræadæ.






Cyprea.

63




64



64



65



66



63



66

Plasma



Gypsea.



67



68



68



70



69



69



71




71



67

G.B.S. jun. exc.



Cypraea.



72



74



73



74



75



Platy



Cypræa.



77




76



77



78



73



78



79



76




79

G.B.S. jun. etc.



Cypræa.





Cypræa.

81



80



81



82



82



80



83



83



Cypræa

88



89



88



90



91




92



91

PLATE 1



Cypræa.




Cypraea.



97



96



96



98



99



98



100



97



100

1000



Cyprea.



101



102



102



103



99



103



104




101



104

G.B.S. jun. etc.

(12000)



Cypræa.



105



106



106



107



108



108



110



110

Opus



Cypræa.



109



105



105



107



III



III



112



112



109

Συρτα



116



117



116



115



113



115



114



117



114

Opis



Κυπρæα



118



121



121



113



120



120




119



119




117




βυρταα.



124



123




123



125.



126



126



127



Plate 1



Pl. 22



151



152



153



154



155



156



157



158



159

Cypræa



125



125



123



126



126



124



127




127

Cypræa



Cypres



Cypræa



145



145



146



146



141



147



147



148



148

GBS per vice

Cypræa



143



142 *



143



141



144



145




142



142

(Spine)



100

Cypræa



G B S Jun^r exc

Fig. 20



Fig. 19



Fig. 18



Fig. 17



Fig. 16



Fig. 15



Fig. 14



Fig. 13



Fig. 12



Fig. 11

Cypræa



125



125



123



126



126



124



127



127

Cypræa



128



128



123



129



129



124



130



130

Cypria



Cypræa



145



145



146



146



141



147



147



148



148

G. B. f. v. c.

Cypræa



143



142 *



143



141



144



144



142



142

Spines



Spines

Cypræa



150



149



149



152



151



151



150

G B S Jun^r exc

Capitulum



281 p. m.

Cypræa



154



153



154



152



153




153



153

G. B. S. jun^r ex



Cyprea



157



158



158



156



160



160

L.B.P. pin. etc.





161



159



161



158



162



162



157



Cypræa

165



163



165





164



167



Cyprina



Cypraea



166



163




165



164



168





Gypræa.



172



172



169



170



170



1



2



3



4



5

Cypræa.




173



174



173



169



171





171



174


G.B.S. jun. acc.



Cypræa.



175




176

G.B.S. jun. exc.




Cypræa.



175




176




(Lobster)





G.B.S. f. m. ecc.



The following Species and Varieties of the Genus

CONUS

Are contained in parts 24, 25, 28, 29, 32, 33, 36, 37, 54 to 57,
147, 148, 151 to 158, of the

CONCHOLOGICAL ILLUSTRATIONS.

FIG.

- 1, 1*. *C. muriculatus* *Nob.* Z. P. 1841. Isl. Masbate, Philippines, on Coral Reefs.
2. *C. albo-maculatus*, *Nob.* Z. P. 1841. (*C. bicolor*. Con. Illus.)
- 3, 3*. *C. maculosus* & var. *Nob.* Z. P. 1841. Isl. Capul. Philippines.
4. *C. Zebra*, *Lam.*
- 5, 5*. *C. Fabula*, *Nob.*
6. *C. nanus*, *Brod.* Z. P. 1833. Lord Hood's Island.
7. *C. aurora*, *Lam.*
- 8, 8*. *C. luteus*, *Brod.* Annaa.
9. *C. arcuatus*, *Nob.* Z. G. IV. 379. Bay of Montija. Sandy mud, at 15 fathoms.
10. *C. tiaratus*, *Brod.* Z. P. 1833. Galapagos Islands.
11. *C. alveolus*, *Nob.* Z. P. 1841.
12. *C. legatus*, *Lam.* Musivum, *Brod.* Z. P. 1833. Philippines.
13. 13*. *C. purpurascens*, *Brod.* Z. P. 1833. Annaa.
14. *C. venulatus*, *Lam.* var. *C. nivifer*, *Brod.* Z. P. 1833. Cape De Verd Islands.
15. *C. planicostatus*, *Nob.* Z. P. 1841.
16. *C. aspersus*, *Nob.* Z. P. 1841. St. Croix. W. J.
17. *C. acuminatus*, var. (*C. insignis*, Con. Illus.)
18. *C. bicolor*, *Nob.* Z. P. 1841.
19. *C. modestus*, *Nob.* perhaps a variety of *cinereus*.
20. *C. unicolor*, *Nob.*
21. *C. vittatus*, *Lam.* var.
22. *C. cardinalis*, *Lam.* var.? (*C. ornatus*, Con. Illus.)

FIG.

23. *C. maculiferus*, *Nob. Z. P.* 1841.
24. *C. bullatus*, *Lam.* var.
25. *C. tornatus*, *Brod. Z. P.* 1833, Panama.
26. *C. geographus*, var. (*C. obscurus*, *Con. Illus.*)
27. *C. Daucus*, *Brug.* var. "lutea, fasciata et punctata."
(*Crocens*, list.)
28. *C. discrepans*.
29. *C. regularis*, var. *Nob. Z. P.* 1841.
30. *C. Princeps*, var. *Brod. Z. P.* 1833. St. Elena.
- 30, a, b, *C. Princeps*, varietates. Monte Christe. Panama.
31. *C. Nux*, *Brod. Z. P.* 1833. Galapagos Isl.
32. *C. concinnus*, *Brod. Z. P.* 1833. Gulph of California.
33. *C. geographus*, var. *rosea*, *Brod. Z. P.* 1833. Annaa.
34. *C. Gladiator*, *Brod. Z. P.* 1833. Panama.
35. *C. artoptus*, *Nob. Z. P.* 1841. S. Seas.
36. *C. incurvus*, *Brod. Z. P.* 1833. Monte Christe.
37. *C. monilifer*, *Sow. Z. P.* 1833. Salango.
38. *C. Archon*, *Brod. Z. P.* 1833. Bay of Montija.
39. *C. Glans*, var. ?
40. *C. Orion*. *Brod. Z. P.* 1833. Rio. Llejos
41. *C. inflatus*, *Nob.*
42. *C. consors*, *Nob. Z. P.* 1841.
43. *C. cinereus*, var.
44. *C. Spectrum*, var.
45. *C. regularis*, *Nob. Z. P.* 1841.
46. *C. Ammiralis*, var. *coronatus*.
47. *C. venulatus*, *Lam.* var. (*C. nivifer*, *Brod.*) Cape De
Verd Isl.
48. *C. Mercator* var. ?
49. *C. Catus*, var.
50. *C. Hyæna* *Lam.* ?
51. *C. sphacelatus*, *Nob. Z. P.* 1841.
52. *C. cedo-nulli*, var. ?
53. *C. splendidulus*, *Nob. Z. P.* 1841.
54. *C. promethæus*, young ? (*C. papillaris* *C. I. list.*)
55. *C. proteus* *Lam.* } var. ? (*C. Breviculus* *C. I. list.*)
Leoninus *Lam.* }
56. *C. Sinensis*. (not fine.) China.

FIG.

57. *C. dispar*
58. *C. balteatus*, *Nob.* Probably a variety of *C. lividus*.
59. *C. concolor*, *Nob.* Z. P. 1841.
60. *C. Mediterraneus*, var, (*C. glaucescens* C. I. list.)
61. *C. pulchellus*, *Nob.* Z. P. 1834.
62. *C. Nussatella*, var.
63. *C. brunneus*, *Wood*, Suppl. pl. 3. f. 1. variety from Galapagos Isl.
64. *C. tendineus*, var. *Annaa*.
65. *C. Elventinus*, *Duclos*, *Guerin*. Icon. Sept. 1833.
66. *C. algoensis*, *Nob.* Z. P. 1834. Algoa Bay.
67. *C. vitulinus*, var.
68. *C. flavescens*, *Gray*.
69. *C. pusio*. *Nob.*
70. *C. textile*, var.
71. *C. auratus* *Lam.* (*C. aulicus*, var ?)
72. *C. nimbosus*, var.
73. *C. monachus*, var ?
74. *C. Capitaneus*, var.
75. *C. flammeus*, *Lam.* var ?
76. *C. solidus*, *Nob.* Z. P. 1841. (Textile, var. list.)
77. *C. mustelinus*, test. jun. ?
78. *C. pauperculus*, *Nob.* Z. P. 1841.
79. *C. maculosus*, test. jun ? (*C. papilliferus* C. I. list.)
80. *C. Thalassiararchus*, *Gray*.
81. *C. miliaris*, var. *granulatus*. (*C. minimus*, C. I. list.)
82. *C. fulgetrum* *Nob.* Z. P. 1841.
83. *C. lividus* *Lam.* var. (*C. Roseus*, *Lam.* C. I. list.)
84. *C. venulatus*, *Lam.* var. (*C. nivifer*, *Brod.* C. I. list.)
85. *C. Thalassiararchus*, var.
86. *C. reticulatus*, *Wood*.
87. *C. Regalitatis*, *Nob.* Z. P. 1834. Real Llejos.
88. *C. brunneus*, *Wood*. (*C. Diadema*, C. I. list.)
89. *C. monachus*, var.
- 90, 90*. *C. Proteus*, test. jun. (*C. Papilionaceus*, test. jun. (Con. Illus. list.)
91. *C. luzonicus*. *Lam* ?
- 92 93. *C. Tulipa*.
94. *C. Cervus*.

- FIG.
 95. *C. geographus*.
 96. *C. bullatus*.
 97. *C. tessellatus*.
 98. *C. tessellatus*, var.
 99. *C. Genuanus*.
 100. *C. Miles*.
 101. *C. eburneus*.
 102. *C. Quercinus*.
 103. *C. Centurio*.
 104. *C. Sumatrensis*.
 105. *C. Barbadosensis*.
 106. *C. Colubrinus*.
 107. *C. Guinaicus*.
 108. *C. cingulatus*, *Tank. Cat. Adamsonii. Gray.*
 109. *C. achatinus*.
 110. *C. aurisiacus*.
 111. *C. Præfectus*.
 112. *C. floccatus*, *Nob. Z. P. 1841. (C. Dux C. I. list.)*
 This shell is much broader, less distinctly striated,
 and more solid than the true *C. Dux*. of Lamarck
 and it is sprinkled with flakes of white.
 113. *C. Generalis, Lam.*
 114. *C. maldivus, Lam.*
 115. *C. Omaicus*.
 116 & 117. *C. Nobilis & var.*
 118, 119. *C. Monile*.
 120, 120*. *C. marmoreus*.
 121. *C. Bandanus, Lam.*
 122, 123. *C. nocturnus, Lam.*
 124. *C. Nicobaricus, Lam.*
 125, 126. *C. araneosus, Lam.*
 127. *C. Zonatus, Lam.*
 128, 129. *C. Imperialis, Lam.*
 130, 131. *C. fuscatus, Lam.*
 132. *C. viridulus, Lam.*
 133, 4, 5. *C. Capitaneus*.
 136. *C. Mustelinus*.
 137. *C. Rattus*

Conus.



1



2



1'



3



4



5



5'



7



6



3'

100. *C. ...*
 101. *C. ...*
 102. *C. ...*
 103. *C. ...*
 104. *C. ...*
 105. *C. ...*
 106. *C. ...*
 107. *C. ...*
 108. *C. ...*
 109. *C. ...*
 110. *C. ...*
 111. *C. ...*
 112. *C. ...*
 113. *C. ...*
 114. *C. ...*
 115. *C. ...*
 116. *C. ...*
 117. *C. ...*
 118. *C. ...*
 119. *C. ...*
 120. *C. ...*
 121. *C. ...*
 122. *C. ...*
 123. *C. ...*
 124. *C. ...*
 125. *C. ...*
 126. *C. ...*
 127. *C. ...*
 128. *C. ...*
 129. *C. ...*
 130. *C. ...*
 131. *C. ...*
 132. *C. ...*
 133. *C. ...*
 134. *C. ...*
 135. *C. ...*
 136. *C. ...*
 137. *C. ...*
 138. *C. ...*
 139. *C. ...*
 140. *C. ...*
 141. *C. ...*
 142. *C. ...*
 143. *C. ...*
 144. *C. ...*
 145. *C. ...*
 146. *C. ...*
 147. *C. ...*
 148. *C. ...*
 149. *C. ...*
 150. *C. ...*
 151. *C. ...*
 152. *C. ...*
 153. *C. ...*
 154. *C. ...*
 155. *C. ...*
 156. *C. ...*
 157. *C. ...*
 158. *C. ...*
 159. *C. ...*
 160. *C. ...*
 161. *C. ...*
 162. *C. ...*
 163. *C. ...*
 164. *C. ...*
 165. *C. ...*
 166. *C. ...*
 167. *C. ...*
 168. *C. ...*
 169. *C. ...*
 170. *C. ...*
 171. *C. ...*
 172. *C. ...*
 173. *C. ...*
 174. *C. ...*
 175. *C. ...*
 176. *C. ...*
 177. *C. ...*
 178. *C. ...*
 179. *C. ...*
 180. *C. ...*
 181. *C. ...*
 182. *C. ...*
 183. *C. ...*
 184. *C. ...*
 185. *C. ...*
 186. *C. ...*
 187. *C. ...*
 188. *C. ...*
 189. *C. ...*
 190. *C. ...*
 191. *C. ...*
 192. *C. ...*
 193. *C. ...*
 194. *C. ...*
 195. *C. ...*
 196. *C. ...*
 197. *C. ...*
 198. *C. ...*
 199. *C. ...*
 200. *C. ...*

Conus.

8



9



8*



10



13



14



11



13*



12

G.B.S. jun. exc.

Conus.

15



15



16



17



18



19




20



21

G.B.S. jun. exc.




Conus

22



23



24



25



26




27



28



29



Conus.



31



30b



32



30b



30a



30a



30



33

G.B.S. jun. exc.




Conus.




35



34



36



37



38



39




40



41



34



Conus.



43



44



42



45



46



48



47



49

G.B.S. jun. exc.



Conus.



51



50



52



53



54



55



56



57



58

G.B.S. jun. exc.

Conus

60



59



61



62



63



64





65



66



67



Conus.

68



69



73



71



72




75



74



70



Conus.

77



76



78



70



80



81



82



83



84



Conus.



85



87



86



88



89



91




90



90*

G.B.S. jun. etc.





93



94



95



92




96



Conus



98



99



97



100




102



101

1847

1847



Conus

103



103



104



106



105



181



182



183



Conus



108



107



109



111



110



Conus

112



112



113



114




114



Conus

117



116



115



119




118






123



121



120




126



Conus



124



125



120*



127



122



Conus

131



133





128



134




135



Conus

130




132



129




131



136





Sowerby, George Brettingham,

The Conchological Illustrations

Lond. 1841

Zool. 498 xe

urn:nbn:de:bvb:12-bsb10308925-4